

THE INFLUENCE OF DIETARY COMPOSITION TO METABOLIC SYNDROME INCIDENCE

by Widhya Aligita

Submission date: 04-Oct-2022 08:10AM (UTC-0500)

Submission ID: 1916378838

File name: 46-Vol.-10-Issue-10-Oct-2019-IJPSR-RA-11617.pdf (561.97K)

Word count: 3228

Character count: 17005



Received on 02 February 2019; received in revised form, 08 July 2019; accepted, 03 September 2019; published 01 October 2019

THE INFLUENCE OF DIETARY COMPOSITION TO METABOLIC SYNDROME INCIDENCE

Widhya Aligita^{*}, Akbar Sumanjaya and Ika Kurnia Sukmawati

Department of Pharmacology, Bandung School of Pharmacy, Jl. Soekarno Hatta 754, Bandung, Indonesia.

Keywords:

Blood glucose,
Blood pressure, HDL,
Metabolicsyndrome, Triglyceride

Correspondence to Author:

Widhya Aligita

Department of Pharmacology,
Bandung School of Pharmacy, Jl.
Soekarno Hatta 754, Bandung,
Indonesia.

E-mail: widhya.aligita@stfb.ac.id

ABSTRACT: Metabolic syndrome is a group of several clinical conditions which is related to the prevalence of another disease such as cardiovascular disease, diabetes mellitus (DM) or increased fasting blood glucose level, visceral obesity, hypertension, *etc.* The major factor that affects the metabolic syndrome occurrence is diet composition. This research aimed to determine the influence of diet composition on metabolic syndrome prevalence. This study was performed using *in-vivo* method. The animal was divided into three groups, each administered a different composition of diet, which was high carbohydrate, high lipid, and high protein diet for 42 days. The parameters were body weight, HDL and triglyceride level, blood glucose level, and blood pressure. The results showed that each diet composition affected the glucose, HDL, triglyceride, and blood pressure level differently. The high-fat diet gave the most significant effect in HDL reducing and blood pressure elevation, while the high carbohydrate diet gave the most significant effect in blood glucose and triglyceride level elevation. It could be concluded that the lowest affecting food to metabolic syndrome was the high protein group, while the high fat and high carbohydrate food had higher risk to induce the metabolic syndrome.

INTRODUCTION: Metabolic syndrome (MetS) is a term for a group of certain clinical conditions which can increase cardiovascular disease risk that is diabetes mellitus or increasing in fasting blood glucose level, visceral obesity, dyslipidemia, and hypertension. There are 20-25% of the world's adult population have metabolic syndrome and those with metabolic syndrome are twice as likely to die from and three times as likely to have a heart attack and stroke compared to those without metabolic syndrome.

In addition, metabolic syndrome raises the fivefold greater risk of diabetes mellitus¹. MetS is a chronic low-grade inflammation as a result of complex interaction between genetic and environmental factor. Several factors affect the development of MetS are visceral adiposity, insulin resistance, endothelial dysfunction, genetic susceptibility, atherogenic dyslipidemia, elevated blood pressure, hypercoagulable state, and chronic stress². Metabolic syndrome is characterized by at least three of the five criteria according to WHO, IDF and NCEP-ATPIII (the National Cholesterol Education Program - Adult Treatment Panel III), that are central obesity (obesity with abdominal circumference that exceeds 80 cm in women and 90 cm in men), elevated triglyceride levels, decreased HDL-cholesterol, fasting glucose up to 110-126 mg/mL (due to decreased insulin sensitivity), and elevated blood pressure^{3,4,5}.

	<p>DOI: 10.13040/IJPSR.0975-8232.10(10).4782-86</p>
	<p>This article can be accessed online on www.ijpsr.com</p>
<p>DOI link: http://dx.doi.org/10.13040/IJPSR.0975-8232.10(10).4782-86</p>	

Metabolic syndrome was said more related to central obesity compared with general obesity. Central obesity was thought to be the main cause of insulin resistance, which was one of the main markers of metabolic syndrome. Furthermore, central obesity could also trigger the release of pro-inflammatory compounds responsible for the diabetes mellitus, hyperlipidemia, and cardiovascular disease occurrence⁶.

“Let food be thy medicine and medicine be thy food”, a phrase from Hippocrates the Ancient Greek father of modern medicine, was indicated that the idea of a relationship between food and health was not a new one. Carbohydrate, lipid, and protein were the primary nutrients or macronutrients we needed. The difference between these three macronutrients was the total amount of energy. The fat contained 9 kcal/g, while protein and carbohydrate contained 54 kcal/g⁷. These differences would determine the total energy intake per day and thus affected the energy balance that in the end could lead to obesity and metabolic syndrome. For example, a high-fat diet could induce weight gain because of excessive energy intake⁸. On the other hand, carbohydrate was also blamed for the incidence of obesity and metabolic syndrome⁹, although it might be because of the wrong choice of the type of carbohydrate^{9,10}. The aim of this study was to evaluate the influence of dietary composition on metabolic syndrome incidence.

MATERIALS AND METHODS:

Animals: Male Wistar mouse 2-3 months old weighing 200-250 g. The animals were kept at standard laboratory conditions at 24-26 °C, humidity 70-75%, and 12 h light/dark cycle. Animals were fed with a treatment diet and water *ad libitum*. The methods in this study were performed in accordance with ethics and guided for animals care and used (139/UN6.KEP/EC/2018).

Administration of Different type Diet Composition in Order to Induce Metabolic Syndrome: Each animals group was fed with 3 different types of diet composition; group 1 was fed with high-fat diet (60% fat), group 2 with high carbohydrate diet (60% carbohydrate), and group 3 with high protein diet (60% protein) for 42 days¹¹. The parameters were body weight, insulin

sensitivity, blood HDL and triglyceride level, and blood pressure. These parameters were measured at before treatment and at 21, 35, and 42 days during treatment.

Insulin Tolerance Test: Before the study was conducted and at 21, 35, and 42 days after treatment, an insulin tolerance test (ITT) using insulin 0.75 U/kg bw, ip, was performed after 4 hours fasting period. Blood glucose concentration from the tail vein was measured using the Easy Touch® blood glucose meter at 0, 15, 30, 45, and 60 minutes after insulin administration¹².

RESULTS:

Effect of Dietary Patterns to Body Weight: The body weight elevation after 42 days administration of different type of diet was shown in **Table 1**.

TABLE 1: BODY WEIGHT PROFILE DURING 42 DAYS OF DIET ADMINISTRATION

Diet composition	Bodyweight at day- (g)			
	0	21	35	42
High fat	208±9	240±19	241±18	237±13
High protein	197±10	234±26	241±14	241±23
High carbohydrate	212±15	250±16	251±14	256±18

Note: Data are presented as mean ± SD, n=6 mice/group, * means significantly different from high protein group, # means significantly different from high carbohydrate group, p<0.05.

Table 1 showed that variation in diet composition, statistically, didn't affect body weight significantly. Although the result showed that high carbohydrate diet gave higher body weight elevation compared to the other groups. And high-fat diet had the lowest effect on body weight.

Effect of Dietary Patterns to HDL Level: The level of HDL during 42 days diet administration was shown in **Table 2**.

TABLE 2: HDL LEVEL DURING 42 DAYS OF DIET ADMINISTRATION

Diet composition	HDL level at day- (mg/dl)			
	0	21	35	42
High fat	24.5 ±5.4	5.3 ±1.5*#	5.2 ±0.7*#	4.5 ±1.2*#
High protein	31 ±11.7	32.6 ±8#	34.5 ±9.6#	35.3 ±8.7#
High carbohydrate	30.7 ±9.1	4 ±0.9*	7.5 ±0.9*	10.3 ±0.9*

Note: Data are presented as mean ±SD, n=6 mice/group, * means significantly different from high protein group, # means significantly different from high carbohydrate group, p<0.05.

From **Table 2**, it could be seen that there were differences between groups treated with high fat, high protein, and high carbohydrate diet. The lowest level of HDL was shown by the high-fat group, followed by carbohydrate group, and the highest level of HDL was shown by high protein group. The HDL level was significantly different compared to each other.

Effect of Dietary Patterns to Triglyceride Level: The level of triglyceride during 42 days diet administration was shown in **Table 3**.

TABLE 3: TRIGLYCERIDE LEVEL DURING 42 DAYS OF DIET ADMINISTRATION

Diet composition	Triglyceride level at day- (mg/dl)			
	0	21	35	42
High fat	64.1 ±20.2	106.3 ±7.2*#	92.1 ±40.5*#	103.8 ±4.6*#
High protein	81.4 ±2.5	81.2 ±4.9#	78.6 ±4.0#	82.1 ±1.3#
High carbohydrate	86.3 ±6.1	124.9 ±33.1*	141.3 ±38.8*	130.3 ±32.4*

Note: Data are presented as mean ±SD, n=6 mice/group, * means significantly different from high protein group, # means significantly different from high carbohydrate group, p<0.05.

Table 3 showed that the triglyceride level was also affected by the different composition of the diet. Different from the HDL level, the highest elevation of triglyceride level was shown by the high

TABLE 5: BLOOD PRESSURE DURING 42 DAYS DIET ADMINISTRATION

Diet composition	Sistol at day- (mmHg)				Diastol at day- (mmHg)			
	0	21	35	42	0	21	35	42
High fat	92±2	114±37*	122±29*	128±24	75±2	85±37	104±21*	109±18*
High protein	97±15	85±2	96±15	103±18	80±16	70±5	77±11	87±17
High carbohydrate	91±3	89±5	99±11	108±20	75±10	72±6	83±13	93±14

Note: Data are presented as mean ±SD, n=6 mice/group, * means significantly different from high protein group, # means significantly different from high carbohydrate group, p<0.05.

Table 5 showed that a high fat diet could significantly increase systolic and diastolic blood pressure, while the high protein and high carbohydrate diet didn't.

DISCUSSION: Body weight was one of the main parameters for metabolic syndrome¹. But from our study, the different compositions of a diet didn't affect the bodyweight significantly. These data were comparable to another study that showed that dietary intake and physical activity could lead to an improvement in metabolic profile, with or without weight loss¹³. From this study, we could see that body weight could not be used as an absolute mark whether a person had metabolic syndrome or not.

carbohydrate group, followed by high fat group, and the lowest triglyceride level was shown by high protein group.

Effect of Dietary Patterns to Fasting Blood Glucose Level: The level of fasting blood glucose during 42 days diet administration was shown in **Table 4**.

TABLE 4: FASTING BLOOD GLUCOSE LEVEL DURING 42 DAYS OF DIET ADMINISTRATION

Diet composition	Fasting blood glucose level at day- (mg/dl)			
	0	21	35	42
High fat	88±15	106±27	124±17*	128±4*
High protein	82±14	86±19	98±25	107±15
High carbohydrate	80±6	129±26*	135±20*	129±13*

Note: Data are presented as mean ±SD, n=6 mice/group, * means significantly different from high protein group, # means significantly different from high carbohydrate group, p<0.05.

Table 4 showed a significant difference between high fat and high carbohydrate group compared to high protein group. But there was no significant difference between high-fat diet compared to high carbohydrate diet.

Effect of Dietary Patterns to Blood Pressure: The level of fasting blood glucose during 42 days of diet administration was shown in **Table 5**.

The HDL level was most affected by the high-fat diet compared to the high carbohydrate and high protein diet. HDL is a lipoprotein that is responsible for transporting lipid from peripheral tissue to liver; in other words it responsible for anti-atherogenic activity¹⁴. The higher the HDL level, the lower the atherogenic risk. Low HDL level, along with high LDL level, could lead to coronary heart disease^{15, 16} and cerebrovascular disease¹⁷. The trans-fatty acid was strongly correlated to the reduction in HDL level¹⁴.

As for the effect on the triglyceride level, the high carbohydrate diet showed the highest elevation compared to the high fat and high protein.

Elevation in triglyceride level was directly affected by the elevation of blood glucose level (hyperglycemia) due to high carbohydrate intake. Hyperglycemia would trigger an increase in insulin requirements resulting in hyperinsulinemia. This condition would eventually cause insulin resistance. Insulin resistance would cause a decrease in the activity of the lipoprotein lipase enzyme, the enzyme responsible for triglyceride transportation to peripheral tissue. The lower the activity of the lipoprotein lipase enzyme, the higher triglyceride level in blood. Then with elevation of triglyceride level, the HDL catabolism would also increase, thus causing a decrease in HDL level^{18, 19}. It was proved by the elevation in HDL level of the high carbohydrate group.

According to the blood glucose level, the high fat and high carbohydrate diet were both had the potential to increase blood glucose level compared to the high protein diet. The mechanism of both causing an elevation in blood glucose level was not fully understood, although it was thought by affecting the insulin sensitivity^{20, 21}.

The high-fat diet was also increasing the blood pressure, while the others didn't. Although, the mechanism of fat-induced hypertension was not fully understood, but another study showed the comparable result²².

From the data above, it could be seen that different diet composition could affect different parameters of metabolic syndrome. And the metabolic syndrome symptoms were not always accompanied by weight gain or obesity. Although it might be because of the lack of time in this experiment to induce obesity. But in general, the high-fat diet and high carbohydrate diet were more responsible in metabolic syndrome symptom occurrence, while the high protein diet gave the lowest effect in metabolic syndrome.

CONCLUSION: From this experiment, it could be concluded that the lowest affecting food to metabolic syndrome was the high protein group, while the high fat and high carbohydrate food had a higher risk in induce the metabolic syndrome.

ACKNOWLEDGEMENT: The authors would like to thank the Bandung School of Pharmacy for permitting to conduct the research work and

providing necessary infrastructure facilities. There are no conflicts of interest for this research work. The authors have not received any funding or research grant for the conduct of this study.

CONFLICT OF INTEREST: The author(s) Widhya Aligita, Akbar Sumanjaya, and Ika Kurnia Sukmawati declare(s) that there is no conflict of interest regarding the publication of this article "The Influence of Dietary Composition to Metabolic Syndrome Incidence".

REFERENCES:

1. International Diabetes Federation: The IDF consensus worldwide definition of the metabolic syndrome 2006. From <http://www.idf.org/metabolic-syndrome>. [Accessed: November 14 2017].
2. Kaur J: A comprehensive review on metabolic syndrome. *Cardiology Research and Practice* 2004; 943162: 1-21. <http://dx.doi.org/10.1155/2014/943162>.
3. NCEP ATP III: ATP III Guidelines At-A-Glance Quick Desk Reference 2001. From <https://www.nhlbi.nih.gov/files/docs/guidelines/atglance.pdf> [Accessed November 12 2017].
4. World Health Organization (WHO): Obesity and Overweight 2017. From <http://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>. [Accessed November 12 2017]
5. International Diabetes Federation (IDF): Diabetes Atlas 8th ed 2017. [ONLINE]. From file:///C:/Users/X441NA/Downloads/english-6th%20(1).pdf [Retrieved November 14 2017].
6. Paley CA and Johnson MI: Abdominal Obesity and Metabolic Syndrome: Exercise as Medicine? *BMC Sports Sci Med Rehabil* 2018; 10: 7.
7. Dam RMV and Seidell JC: Carbohydrate Intake and Obesity. *European Journal of Clinical Nutrition* 2007; 61 (S 1): S75-S99.
8. Bray GA, Paeratakul S and Popkin BM: Dietary fat and obesity: a review of animal, clinical and epidemiological studies. *Physiol Behav* 2004; 83: 549-55.
9. Pitsavos C, Panagiotakos D, Weinem M and Stefanadis C: Diet, Exercise, and the Metabolic Syndrome. *The Review of Diabetic Studies* 2006; 3(3): 118-26.
10. Di-Meglio DP and Mattes RD: Liquid versus Solid Carbohydrate: Effects on Food Intake and Body Weight. *International Journal of Obesity* 2000; 24: 794-00.
11. Aligita W, Kurniati NF and Sukandar EY: Antidiabetic Study of Combination of *Andrographis paniculata* (Burm. F.) Wallich. Ex Nees. herbs extract and *Guazuma ulmifolia* Lamk. leaves extract in obese diabetic mice model. *Int J Pharm Pharm Sci* 2016; 8: 316-20.
12. Aligita W, Susilawati E, Sukmawati IK, Holidayanti L and Riswanto J: Antidiabetic activity of *Muntingia calabura* L. leaves water extract in type 2 diabetes mellitus animal model. *The Indonesian Biomedical Journal* 2018; 10(2): 165-70.
13. Keane D, Kelly S, Healy NP, McArdle MA, Holohan K and Roche HM: Diet and Metabolic Syndrome: An Overview. *Current Vascular Pharmacology* 2013; 11: 1-16.
14. Yanai H, Katsuyama H, Hamasaki H, Abe S, Tada N and Sako A: Effect of dietary fat intake on HDL metabolism. *Journal of Clinical of Medicine Res* 2015; 7(3): 145-49.

15. Kitamura A, Iso H, Naito Y, Iida M, Konishi M, Folsom AR, Sato S, Kiyama M, Nakamura M, Sankai T, Shimamoto T and Komachi Y: High-density lipoprotein cholesterol and premature coronary heart disease in urban Japanese men. *Circulation* 1994; 89(6): 2533-39.
16. Yokokawa H, Yasumura S, Tanno K, Ohsawa M, Onoda T, Itai K, Sakata K, Kawamura K, Tanaka F, Yoshida Y, Nakamura M, Terayama Y, Ogawa A and Okayama A: Serum low-density lipoprotein to high-density lipoprotein ratio as a predictor of future acute myocardial infarction among men in a 2.7-year cohort study of a Japanese northern rural population. *J Atheroscler Thromb* 2011; 18(2): 89-98.
17. Okamura T, Hayakawa T, Kadowaki T, Kita Y, Okayama A and Ueshima H: The inverse relationship between serum high-density lipoprotein cholesterol level and all-cause mortality in a 9.6-year follow-up study in the Japanese general population. *Atherosclerosis* 2006; 184(1): 143-50.
18. Mohammed NH and Wolever TM: Effect of carbohydrate source on post-prandial blood glucose in subjects with type 1 diabetes treated with insulin lispro. *Diabetes Res Clin Pract* 2004; 65: 29-35.
19. Wolever TM and Mehling C: Long-term effect of varying the source or amount of dietary carbohydrate on postprandial plasma glucose, insulin, triacylglycerol, and free fatty acid concentrations in subjects with impaired glucose tolerance. *Am J Clin Nutr* 2003; 77: 612-21.
20. Hu FB, Van Dam RM and Liu S: Diet and risk of type ii diabetes: the role of types of fats and carbohydrates. *Diabetologia* 2001; 44: 805-17.
21. Marshall JA and Bessesen DH: Dietary fat and the development of type 2 diabetes. *Diabetes Care* 2002; 25(3): 620-22.
22. Burstyn PG and Husbands DS: Fat induced hypertension in rabbits: effects of dietary fiber on blood pressure and blood lipid concentration. *Card Res* 1980; 14(4): 185-91.

How to cite this article:

Aligita W, Sumanjaya A and Sukmawati IK: The influence of dietary composition to metabolic syndrome incidence. *Int J Pharm Sci & Res* 2019; 10(10): 4782-86. doi: 10.13040/IJPSR.0975-8232.10(10).4782-86.

All © 2013 are reserved by International Journal of Pharmaceutical Sciences and Research. This Journal licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License.

This article can be downloaded to **ANDROID OS** based mobile. Scan QR Code using Code/Bar Scanner from your mobile. (Scanners are available on Google Playstore)

THE INFLUENCE OF DIETARY COMPOSITION TO METABOLIC SYNDROME INCIDENCE

ORIGINALITY REPORT

24%

SIMILARITY INDEX

16%

INTERNET SOURCES

16%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

1	rasayanjournal.co.in Internet Source	4%
2	innovareacademics.in Internet Source	3%
3	Submitted to Universiti Malaysia Pahang Student Paper	3%
4	Wesley Wong, Austin Kim, Ashley W. Seifert, Malcolm Maden, Justin D. Crane. " Spiny mice () exhibit attenuated hallmarks of aging and rapid cell turnover after UV exposure in the skin epidermis ", Cold Spring Harbor Laboratory, 2020 Publication	2%
5	strathprints.strath.ac.uk Internet Source	1%
6	valleyinternational.net Internet Source	1%
7	Balazs, T.. "The influence of excess body fat on the cardiotoxicity of isoproterenol in rats",	1%

Toxicology and Applied Pharmacology,
196209

Publication

8

Jiao Gong, Jingyi Ou, Xueping Qiu, Yusheng Jie et al. "A Tool to Early Predict Severe Corona Virus Disease 2019 (COVID-19) : A Multicenter Study using the Risk Nomogram in Wuhan and Guangdong, China", Cold Spring Harbor Laboratory, 2020

Publication

1 %

9

cris.maastrichtuniversity.nl

Internet Source

1 %

10

Sophia Rose, Fillah Fithra Dieny, Nuryanto Nuryanto, A. Fahmy Arif Tsani. "The Correlation between Waist-to-Height Ratio (WHtR) and Second to Fourth Digit Ratio (2D:4D) with an Increase in Metabolic Syndrome Scores in Obese Adolescent Girls", Electronic Journal of General Medicine, 2020

Publication

1 %

11

Submitted to Uttar Pradesh Technical University

Student Paper

1 %

12

Deirdre Keane, Stacey Kelly, Niamh Healy, Maeve McArdle, Kieran Holohan, Helen Roche. "Diet and Metabolic Syndrome: An Overview", Current Vascular Pharmacology, 2014

1 %

13	Suljkovicova, H.. "Effect of macronutrient composition of the diet on the regulation of lipolysis in adipose tissue at rest and during exercise: Microdialysis study", Metabolism, 200210 Publication	1 %
14	www.mdpi.com Internet Source	1 %
15	www.nature.com Internet Source	1 %
16	Submitted to University of Wales Institute, Cardiff Student Paper	1 %
17	www.cdc.gov Internet Source	1 %
18	www.inabj.org Internet Source	1 %
19	Submitted to Edge Hill University Student Paper	1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

THE INFLUENCE OF DIETARY COMPOSITION TO METABOLIC SYNDROME INCIDENCE

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
