Rasayan J. Chem. 17/10/22 20.37



About Rasayan J. Chem.



Rasayan J. Chem. ISSN: 0974-1496(Print) ISSN: 0976-0083(Online) CODEN: RJCABP

Frequency: March, June, September & December

The word RASĀYAN is the true inspiration behind the name of this journal, which comes from Sanskrit origin, which means Chemical- is an incredibly reactive and self-explanatory term. Which has been widely used in our scriptures and literature since ancient times.

Established in 2008, RASĀYAN Journal of Chemistry is a peer-reviewed international quarterly research journal of chemical science with its multidisciplinary nature, which is SCOPUS(Elsevier) indexed since 2008 and published as open access as well as a print journal.

RASĀYAN Journal of Chemistry (RJC) widely covers all branches of Chemistry including: Organic, Inorganic, Physical, Analytical, Biological, Pharmaceutical, Industrial, Environmental, Agricultural & Soil, Petroleum, Polymers, Nanotechnology, Green Chemistry, Forensic, Phytochemistry, Synthetic Drugs, Computational, as well as Chemical Physics and Chemical Engineering. In the online version of the journal, all figures and photographs are colored with printable clarity.

- RASĀYAN Journal of Chemistry (RJC) has been included in the consolidated List of journals recently recommended by the UGC.
- Types of manuscripts accepted in RJC: Full-length research papers, Review articles and Short/ Rapid communications.
- Authors may also submit their recently published books for Book Review.
- Apart from its regular issues, RASĀYAN Journal of Chemistry is also welcoming to the proposals of the thematic and special issues on emerging areas of research.
- Editor-in-Chief may also invite prominent scientists/ researchers to write articles on some specific topics related to cutting edge researches.



CONTENTS

ARCHIVE ISSUES

DOWNLOADS

RJC CONFERENCE

SPECIAL ISSUES-



2.1 Cite Score

45th percentile
Powered by Scopus

Rasayan J. Chem. 17/10/22 20.37

Powered By









RJC is Abstracted / Indexed
















This work is licensed under a <u>Creative</u>
<u>Commons Attribution</u>
<u>4.0 International</u>
<u>License.</u>



















Important Links

Home
About us
Contact
Archive
Abstracted
Announcement

Editorial Board
Author's Guidelines
Subscription
Manuscript Submission
Current Issue

About Rasayan

Rasayan J. Chem.
ISSN: 0974-1496(Print)
ISSN: 0976-0083(Online)
CODEN: RJCABP
Frequency: March, June, September &
December

Editorial Board 17/10/22 20.37

Editorial Board

Editor-in-Chief:



Sanjay K. SHARMA, FRSC

Professor, Department of Chemistry & Dean(Research), JECRC University, Jaipur, India

Contact: +91 9001699997

Email: editor@rasayanjournal.com,

Research Interest: Green Chemistry, Organic Chemistry and Water

treatment

Editorial Office:



Pratima SHARMA

Publisher and Managing Ed RASĀYAN Journal of Chemis 23 'Anukampa', Janakpuri, O

Stn., Ajmer Road, Jaipur-302024 (India)

Contact: 9414202678

Email: rasayanjournal@gmail.co

CURRENT ISSUE Online First



Bassim H. Hammadi

Department of Chemical Engineering, College of Engineering, Qatar University, P.O. Box 2713, Doha, Qatar

Contact: +97440434142 Email: b.hammadi@qu.edu.qa

Research Interest: Reaction Engineering, Adsorption Technology



Florent ALLAIS

Director, R&D Unit of Industrial Agro-Biotechnologies URD

ABI- AgroParis Tech, Pomacle, France

Contact: +33 633 698 126

Email: Florent.allais@agroparistech.fr

Research Interest: Green Chemistry, Bio-based Polymers



Goutam BRAHMACHARI

Professor, Chemistry Department, Visva-Bharati University, Santiniketan-731235, India.

Contact: +91 943485744

Email: goutam.brahmachari@visva-bhartai.ac.in

Research Interest: Organic Synthesis; Green Chemistry; Natural

products, Medicinal Chemistry



Ishmael MASESANE

Professor, Department of Chemistry, University of Botswana, Botswana

Contact: 26772874348

Fmail: MASESANE@UB.AC.BW

Research Interest: Organic synthesis, Natural product Chemistry,

Medicinal Chemistry



Marei Mailoud EL-AIAILY

University of Benghazi, Faculty of Science, Department of Chemistry, Benghazi, Libya



Eno E. EBENSO

Professor, North-West University Gauteng, South Africa

Contact: +27825387286

Email: Eno.Ebenso@nwu.ac.za

Research Interest:

DOWNLOADS

RJC CONFERENCE



Giusy LOFRANO

Department of Environment, University of Salerno, Salerno, Italy

Contact: 0039 347 90 60 670 Email: glofrano@unisa.it

Research Interest: nanotechnologies, wa

oxidation processes

SCOPUS ed Since, 2008

Approved

UGC

ced



Hakan ARSLAN

Department of Chemistry, Faculty of Arts and Science, Mersin University, Mersin, TR-33343, Turkey

Contact: +90.532.7073122

Email: hakan.arslan@mersin.edu.tr

CiteScore

Research Interest: Coordination chemist 45th percentile

Kinetic Studies, X-ray diffraction studies, Spectroscopy



Ime Bassey OBOT

Chemistry

Center of Research Excellence in Corrosion Research Institute, King Fahd University of Potroloum and Minorals

(KFUPM), P.O. Box 489, Dhahran, 3 Rasayan Journal of

Contact: +966 13 860-8283

Research Interest: Corrosion and Sca

Computational Chemistry.

Email: obot@kfupm.edu.sa





Man SINGH

Professor and Dean, school of Chemical sciences, Gujrat central University, Gandhinagar, Gujrat, India

Contact: +91 9408635094

Editorial Board 17/10/22 20.37

Contact: 00218918315683
Email: melajaily@gmail.com

Research Interest: Mixed ligand complexes, Drugs, Applications,

Corrosion inhibition, Molecular docking, DFT studies



Mika SILLANPÄÄ

Department of Chemical Engineering, School of Mining, Metallurgy and Chemical Engineering, University of Johannesburg, Doornfontein 2028, South Africa

Contact: +358400205215

Email: mikaesillanpaa@gmail.com

Research Interest: Water treatment



Pankaj KUMAR

Professor and Head, Department of Chemistry, University of Energy and Petrolium studies, Dehradun, India

Contact: +917351958165

Email: pkumar@ddn.upes.ac.in

Research Interest: Biofuels and Bioenergy, Chemical sensors, Nanomaterials, Minimization of industrial wastes



R.V. SINGH

Ex Professor, Department iof Chemistry, University of Rajasthan, Jaipur, India

Contact: +91 941406975
Email: rvsjpr@hotmail.com

Research Interest: Inorganic Chemistry



Soro YAYA

Laboratoire des Procédés Industriels de Synthèse, de l'Environnement et des Energies Nouvelles (LAPISEN), Institut National Polytechnique (INP-HB),Yamoussoukro, BP 991 Yamoussoukro(Côte d'Ivoire)

Contact: (+225) 07 71 67 66 **Email**: <u>soro_y@yahoo.fr</u>

Research Interest: Organic synthesis, Natural Products, waste management



V.K. GARG

Professor and Dean Centre for Environmental Science and Technology School of Environment and Earth Sciences Central University of Punjab, Bathinda- 151001, India

Contact: +919812058109
Email: vinodkgarg@yahoo.com

Research Interest: Pollution Monitoring and abatement, Solid Waste

Management, Radioecology

Email: mansingh50@hotmail.com

Research Interest: Surface Chemistry, Pl





Nnabuk Okon EDDY

This work is licensed under a <u>Creative</u>

12,293 Pageviews

Sep 17th

Oct 17th

Professor, Department of Chemistry, Allinadu Bello University, Zaria, Kaduna State, Nigeria

Contact: +2348038198753

Email: nabukeddy@yahoo.com

Research Interest: Physical Chemistry, Readletts a councily the Nanochemistry, Industrial Chemistry, Environmental Westerdy



Priyan PERERA

Director - Center for Sustainability
Department of Forestry and Enviro
University of Sri Jayewardenepura

Contact: +94 718656457

Email: priyan@sjp.ac.lk

Research Interest: Environmental Sustainability, Ecology



Ramesh. L. GARDAS

Department of Chemistry Indian Institute of Technology Madras Chennai-600 036, India

Contact: +91 9884996125

Email: _gardas@iitm.ac.in

Research Interest: Physical Chemistry, Chemical Thermodynamics,

Alternative Solvents



Susheel MITTAL

Senior Professor, School of Chemistry & Biochemistry, Thapar Institute of Engineering & Technology (Deemed to be University), Bhadson Road, Patiala-147004, India

Contact: +91-9815653261

Email: smittal2001@yahoo.com

Research Interest: Voltammetric Sensors, Potentiometric Sensors,

Biosensors, Ambient Air Quality and Human Health



Willian Aperador CHAPARRO

School of Engineering, Universidad Militar Nueva Granada, Bogotá-111121, Colombia

Contact: + 57 3142220552

Email: william.aperador@unimilitar.edu.co

Research Interest: Materials, batteries, corrosion, coatings, tribology

Editorial Board 17/10/22 20.37

Important Links

Home

About us

Contact

Archive

Abstracted

Announcement

Editorial Board
Author's Guidelines
Subscription
Manuscript Submission

Current Issue

About Rasayan

Rasayan J. Chem. ISSN: 0974-1496(Print) ISSN: 0976-0083(Online) CODEN: RJCABP

Frequency: March, June, September &

December

© Copyright 2020 RJC. All rights reserved. website designed and managed by ubijournal.



Archive Issue



Volume 13 Number 1, 1-779, January- March (2020)

ARCHIVE ISSUES

DOWNLOADS

THE EFFECTIVENESS OF AQUAPONIC COMPARED TO MODIFIED CONVENTIONAL AQUACULTURE FOR IMPROVED OF

AMMONIA, NITRITE, AND NITRATE

RJC CONFERENCI



<u>- Deswati, Amelliza Deviona , Ella Intan Sari , Yulizar Yusuf and Hilfi Pardi</u>

UPDATES ON 4-HYDROXYDERRICIN AND XANTHOANGELOL OF ANGELICA PLANTS: EXTRACTION AND PHARMACOLOGICAL ACTIVITIES

<u>– D. L. Aulifa, I. K. Adnyana , Sukrasno and J. Levita</u>

SCOPUS

UGC Approved





IDENTIFICATION OF SECONDARY METABOLITE OF LABAN LEAF EXTRACT (VITEX PINNATA L) FROM GEOTH AND NON-GEOTHERMAL OF AGAM MOUNTAINS IN ACEH BESAR, ACEH PROVINCE, INDONESIA

- Cut Nuraskin, Marlina , R. Idroes , C. Soraya and Djufri

A STRAIGHT FORWARD, FAST, SENSITIVE, EPI- HPLCMS/MS METHOD FOR THE SIMULTANEOUS ESTIMATIC CLOZAPINE AND AMITRIPTYLINE IN HUMAN PLASMA

- Muni Krishnaiah A , C. Rama Chandraiah, M. Srinivas and N. Devanna

45th percentile



DYE-SENSITIZED SOLAR CELLS PROPERTIES FROM NATURAL DYE AS LIGHT-REAPING MATERIALS EXTRAG GAYO ARABICA COFFEE HUSKS

<u>- Hasby, Nurhafidhah , Gawang Pamungkas , and Said Ali Akbar</u>

Rasayan Journal of Chemistry

VAPOUR GENERATION CHEMILUMINESCENCE DETECTION SYSTEM FOR DETERMINATION OF <u>DIMETHYLSULPHONIOPROPIONATE AND DIMETHYLSULPHIDE IN TROPICAL SEAWATER</u>

A. E. Adedapo, N. U. Benson, A. B. Williams and K. Toda

Pharmacology, Toxicology and Pharmaceutics... best quartile

SJR 2021 0.33

powered by scimagojr.com

DESIGN OF A NOVEL EQUIPMENT FOR TREATMENT OF GREYWATER USING WOOD APPLE SHELL AS AN

Kalyani Gaddam and Lakshmi Adathodi

 \bullet BIOPROSPECTING CELLULOLYTIC FUNGI ASSOCIATED WITH TEXTILE WASTE AND INVITRO OPTIMIZATION CELLULASE PRODUCTION BY Aspergillus flavus NFCCI-4154 - <u>Priya Sutaoney, Rachana Choudhary and A.K. Gupta</u> This work is licensed under a Creative PREPARATION AND CHARACTERIZATION OF GOATSKIN GELATIN AS HALAL ALTERNATIVE TO BOVINE GELATION OF Attribution 4.0 International License. <u> A. Bahar , N. Kusumawati and S. Muslim</u> CHARACTERISTIC OF HYDROXYAPATITE-SUPPORTED NI, Ce, Cu CATALYST PREPARED BY DEPOSITION PRECIPITATION Readers around the AND IMPREGNATION METHOD FOR GLYCEROLAQUEOUS PHASE REFORMING PROCESS world L. Hakim, H. Husin, R. Sari, R. Dewi and Meriatna 12,298 Pageviews SYNTHESIS, CHARACTERIZATION, SURFACE FUNCTIONALIZATION OF MAGNETITE NANOPARTICLES AND ANTIBACTERIAL STUDIES <u>— Jaiveer Singh, Arti Jangra, Jai Kumar, Keerti Rani and Ramesh Kumar</u> KINETICS AND MECHANISM OF THE AQUEOUS PHASE OXIDATION OF HYDROGEN SULFIDE BY OXYGEN: CATALYZED BY HYDROQUINONE <u> – Deepak Singh Rathore and C. P. Singh Chandel</u> THE MODIFICATION OF CYANIDIN BASED DYES TO IMPROVE THE PERFORMANCE OF DYE-SENSITIZED SOLAR CELLS. (DSSCs) <u>– Imelda , Emriadi, Hermansyah Aziz, Adlis Santoni and Nofitri Utami</u> SYNTHESIS AND ANTIOXIDANT ACTIVITY STUDY OF NEW MANNICH BASES DERIVED FROM VANILLIC ACID <u>- H. Hayun, I. Gavrila, S. Silviana, A.E.K. Siahaan, R.F. Vonna and M.I. Latifah</u> REMOVAL OF CO, SO2 AND HC GASES FROM INCINERATOR USING DIATOMACEOUS EARTH AND COMPOST-MODIFIED SORBENT IN FIXED BED ABSORPTION REACTOR <u>– Mariana, F. Mulana and L. Juniar</u> ASSESSMENT OF WATER QUALITY OF SITNICA RIVER BY USING WATER QUALITY INDEX (WQI) - A. M. Shala Abazi , B. H. Durmishi , F. S. Sallaku , H. S. ‡adraku , O. B. Fetoshi, P. H. Ymeri and P. S. Bytv§i

UNSYMMETRIC COMPARTMENTAL LIGANDS FOR METALLOBIOSITE MODELLING: A MINI REVIEW

– Dipesh Ghosh



PHYSICOCHEMICAL AND BIOLOGICAL WATER QUALITY ASSESSMENT OF THE GUEBLI RIVER, NORTHEASTERN ALGERIA

- <u>K. Boudeffa, F. Fekrache and N. Bouchareb</u>



ISOLATION AND CHARACTERISATION OF CELLULOSE NANOCRYSTAL OBTAINED FROM SUGARCANE PEEL

<u>– C. V. Abiaziem, A. B. williams , A. I. Inegbenebor , C. T. Onwordi , C. O. Ehi-Eromosele and L. F. Petrik</u>



BIOSYNTHESIS OF FUNCTIONALIZED GOLD NANOPARTICLES BY USING METHYL COMMATE C IN Scoparia dulcis LEAF EXTRACT AS REDUCING AGENT



<u> – J. Mary Joselin, V. Ganesh Kumar, T. Selvaraj, K. Govindaraju and V. Karthick</u>

DETERMINATION OF HEAVY METALS IN PRISTIPOMA FURCATUS AND ACANTHURUS STRIGOSUS FISH SPECIES COLLECTED FROM PULICAT LAKE, CHENNAI



– B. Prabhu Dass Batvari and D. Saravanan

POLYANILINE (PANI)-SENSITIZED Fe3O4/SiO2/TiO2 NANOCOMPOSITES AS PHOTOCATALYST FOR THE REDUCTION OF AU(III) IONS



<u>– I. S. Budi, S. J. Santosa and E. S. Kunarti</u>



ASSESSMENT OF TOXIC METALS IN COMMON FOOD GRAINS GROWN IN JODHPUR CITY



<u>— Pallavi Mishra, Rajshri Soni, Vandana Kachhwaha and Naresh Giri</u>

SYNTHESIS, CHARACTERIZATION AND ANTIMICROBIAL ACTIVITY OF Mn(II),Fe(II), Ni(II),Co(II) AND Zn(II) COMPLEXES OF SCHIFF BASE DERIVED FROM 2,2- DIMETHYLPROPANE 1, 3-DIAMINE AND 5-CHLORO ISATIN



<u>— N. P. Singh, K. Kumar , A. Kumar and U. Agarwal</u>

DEGRADATION STUDYOF PIMAVANSERIN: IDENTIFICATION, ISOLATION AND STRUCTURAL CHARACTERIZATION OF DEGRADANTS



— Shaik. John Saida, A. Manikandan , V.V.S.R.N. Anji Karun Mutha , Muralidharan Kaliyaperumal , Chidananda Swamy Rumalla , Ramulu Yanaka and S. Venkat Rao



Mangifera odorata GRIFF SEED EXTRACT AS CORROSION INHIBITOR OF MILD STEEL IN HYDROCHLORIC ACID MEDIUM



<u>— Y. Stiadi, Rahmayeni, L. Rahmawati, M. Efdi, H. Aziz and Emriadi</u>

OPTIMIZATION OF ENCAPSULATION EFFICIENCY AND LACTIC ACID BACTERIA VIABILITY THROUGH A COMBINATION BETWEEN CAPSULE AGENTS AND TOFU WASTE PREBIOTIC



<u> – Virna Muhardina, Putri Meutia Sari , Yuliani Aisyah , Sri Haryani and Said Ali Akbar</u>

THEORETICAL VALIDATION OF MEDICINAL PROPERTIES OF Curcuma longa Linn



- Sampat Suryawanshi and Pramod Kulkarni

SILICA EXTRACTION FROM BEACH SAND FOR DYES REMOVAL: ISOTHERMS, KINETICS AND THERMODYNAMICS



| <u>— M. Lutfi Firdaus, Fitri E. Madina , Sasti Yulia F., Rina Elvia , Soraya N. Ishmah , Diana R. Eddy. Abigail P. Cid-Andres</u>

GROWTH CONTROL OF ZnO-TiO2/CHITOSAN NANOROD ON COTTON TEXTILE FIBER BASED ON DIFFERENT CHLORO ACETIC MOLAR COMPOSITION AS CROSS LINKER



<u> — Yetria Rilda, Dyah Rahayu Ratyaningsih , Yulia Eka Putri , Refinel Syukri , Anthoni Agustien and Hilfi Pardi</u>

STUDY OF ACOUSTICAL PARAMETERS OF MONOMETHYLAMMONIUM PERCHLORATE IN SOME NON-AQUEOUS SOLVENTS AT DIFFERENT TEMPERATURES USING ULTRASONIC TECHNIQUE



<u> – Vivek Pathania , Shrutila Sharma, Shashi Kiran Vermani and B.K.Vermani</u>

MECHANICAL PROPERTIES OF BIOPLASTICS JANENG STARCH (Dioscorea hispida) FILM WITH GLYCEROL AND ZINC OXIDE AS REINFORCEMENT



<u> – Chairul Amni , Ismet , Sri Aprilia , Mariana and Said Ali Akbar</u>

SYNTHESIS OF NEW SULPHANILAMIDE BASED SCHIFF BASE NICKEL COMPLEXES WITH STUDY OF ITS ANTIBACTERIAL ACTIVITY AND NANOPARTICLE SYNTHESIS



<u>– R. R. Surve and S. T. Sankpal</u>

SYNTHESIS OF FEW 1,3,4-OXADIAZOLE DERIVATIVES BLENDED WITH DIFFERENT HETEROCYCLES AND THEIR IN-VITRO ANTIBACTERIAL ACTIVITIES



<u>— M. Idrees, Y.G. Bodkhe , N. J. Siddiqui and S. Kola</u>

SYNTHESIS OF MgFe2O4-MgO NANOCOMPOSITE: INFLUENCE OF MgO ON THE CATALYTIC ACTIVITY OF MAGNESIUM FERRITE IN BIODIESEL PRODUCTION



<u>— Helmiyati , G.H. Abbas, Y. Budiman and S. Ramadhani</u>

PHYTOCHEMICAL GINKGOLIDE B PROTECTS CULTURED NEUROBLASTOMA SH-SY5Y CELLS AGAINST Aβ(25-35) INDUCED OXIDATIVE STRESS RESPONSES BY MAINTAINING THE MITOCHONDRIAL INTEGRITY



<u> – Navrattan Kaur , Sharanjot Kaur , Meenu Saini , Monisha Dhiman and Anil K. Mantha</u>





<u>— Purwantiningsih, N. Jannah and D. U. C. Rahayu</u>

BIOACTIVE COMPOUND FROM THE MANGROVE PLANT ENDOPHYTIC FUNGUS Diaporthe amygdali SgKB4



<u> D. Handayani, T. Wahyuningsih , Rustini , M.A. Artasasta , A. E. Putra and P. Proksch</u>





<u>- Zalak J. Patel, Mukesh C. Patel , Parimal M. Chatrabhuji , Viral A. Patel and Dharmesh R. Patel</u>



DETERMINATION OF THE CONDITIONS OF PHOSPHATE COATINGS FORMATION ON IRON BY VOLTAMMETRIC METHOD



<u>– V. N. Statsyuk, A. Bold, L. A. Fogel, U. Sultanbek and Zh. Tilepbergen</u>

RESULTS OF THE STUDY OF FILMS OBTAINED BY ADDING VARIOUS IMPURITIES TO THE SOLUTION OF NICKEL PHOSPHIDE



<u> P. Abdurazova, Sh. Koshkarbayeva, M. Sataev, N. Dzhakipbekova and Y. Raiymbekov</u>

ELECTROPHORETIC DEPOSITION OF ZIRCONIA ON ALUMINA FOR MOLTEN SALT APPLICATIONS A. Sheik Mideen, G. Venkatasubramanian, T.M. Sridhar and S. Vasudevan SYSTEMATIC LC-MS/MS METHOD FOR QUANTIFICATION OF 2,3-DIMETHYL-2H-INDAZOLE-6-AMINE CONTENT IN PAZOPANIB HYDROCHLORIDE <u>— Pikkili Viswanath, Doddipalli Venkata Ramana Reddy and Nagaraju Chamarthi</u> GREEN ROUTE FOR THE SYNTHESIS OF OXADIAZOLE DERIVATIVE CONTAINING BENZIMIDAZOLE MOIETY AND ITS MANNICH BASES: IN-VITRO ANTIMICROBIAL ACTIVITY <u> – Samrin Naz and Mahendra B. Bagade</u> ANTIMICROBIAL METABOLITE FROM THE ENDOPHYTIC FUNGI Aspergillus flavus ISOLATED FROM Sonneratia alba, A MANGROVE PLANT OF TIMOR-INDONESIA — Antonius R. B. Ola, Christina A. P. Soa, YosephSugi , Theo Da Cunha, Henderiana L. L. Belli and Herianus J. D. Lalel PHYSICOCHEMICAL CHARACTERIZATION OF NATURAL KAOLIN FROM JABOI INDONESIA <u> R. Dewi, H. Agusnar, Z. Alfian and Tamrin</u> THE ENCAPSULATION OF METFORMIN ON CHITOSAN MATRIX AS DIABETES MELLITUS DRUG SLOW RELEASE SYSTEM <u>- S. E.Cahyaningrum , N. Herdyastuti, W. Imroni and A. Sholikhah</u> GROWTH, CHARACTERIZATION OF Mn(II) DOPED Zn(II) L-HISTIDINE HYDROCHLORIDE MONOHYDRATE COMPLEX **CRYSTALS** - V. Parvathi, J. Sai Chandra , Y. Sunandamma and H.Ananda Lakshmi ADSORPTION CORROSION INHIBITIVE BEHAVIOR OF PEELS EXTRACT OF Theobroma cacao ON MILD STEEL AS A CORROSION INHIBITOR IN HCL MEDIA <u>- Y. Yetri, R. Hidayat, R. Sumiati and N. P. Tissos</u> BIOACTIVE METABOLITE FROM MARINE SPONGEDERIVED FUNGUS Cochliobolus geniculatus WR12 <u> – D. Handayani, R. A. Putri , F. Ismed , T. Hertiani N. P. Ariantari and P. Proksch</u> Teedia Lucida ROOT EXTRACTS BY ULTRASONICATION AND MACERATION TECHNIQUES: PHYTOCHEMICAL SCREENING, ANTIMICROBIAL AND ANTIOXIDANT POTENTIALS - P.T. Motsumi , T. Qwebani-Ogunleye , I.P. Ejidike , F.M. Mtunzi and Z. Nate EFFECTIVE METHODS OF PYRIDOXINE SUPPLEMENTATION IN LAYING HENS TO ALBUMIN AND GLOBULIN LEVELS <u>- S. Silaban, B. Sinaga , M. Damanik and P. M. Silitonga</u>

POROSITY EFFECT OF Ag DOPED ZnO NANOCRYSTALLITES

— M. Giruba, J. Christina Rhoda, S. Chellammal and K. Ravichandran

COMPARATIVE STUDY ON ADSORPTION OF CATIONIQUE DYE ONTO EXPANDED PERLITE AND NATURAL CLAY - B. Damiyine, A. Guenbour and R. Boussen IDENTIFICATION, CHARACTERIZATION AND ANTIBACTERIAL POTENTIAL OF PROBIOTIC LACTIC ACID BACTERIA ISOLATED FROM NANIURA (A TRADITIONAL BATAK FERMENTED FOOD FROM CARP) AGAINST Salmonella typhi <u>— G. Haro , I. Iksen, and N. Nasri</u> MOLECULAR DOCKING ANALYSIS OF GOSSYPOL ANALOGUES AS HUMAN NEUTROPHIL ELASTASE (HNE), MATRIX METALLOPROTEINASES (MMP 2 AND 9) AND TYROSINASE INHIBITORS Vishnu Ragavan , Aishwarya Ramesh and Radhakrishnan Narayanaswamy PHYTOCHEMICAL SCREENING AND TOXICITY OF ETHANOLIC EXTRACT OF MANGROVE (Rhizophora mucronata) LEAVES FROM LANGSA, ACEH TIMUR <u>— Gimelliya Saragih, Tamrin , Marpongahtun , Darwin Yunus Nasution and Abdillah</u> WETTING ABILITY OF A PHYTOPREPARATION AND THEIR ASSOCIATES WITH POLYELECTROLYTES. <u> — O. Yessimova , S. Kumargaliyeva , M. Kerimkulova , K. Mussabekov and Zh. Toktarbay</u> NOVEL HETEROPOLYACID SALT: TITANIUM (IV) MOLYBDOTUNGSTATE - <u>N. Sharma, H.K. Sharma and P. Dogra</u> SYNTHESIS OF 4-CHLORO-PIPERIDINE DERIVATIVES VIA NDCL5 MEDIATED AZA-PRINS TYPE CYCLIZATION OF EPOXIDES AND HOMOALLYLIC AMINES - K. Kamesu, G. V. Krishna Mohan and K. Rajasekhar EVALUATION OF TETRA-n-BUTYLAMMONIUM BROMIDE AS CORROSION INHIBITOR FOR MILD STEEL IN 1N HCL MEDIUM: EXPERIMENTAL AND THEORETICAL INVESTIGATIONS <u> N. Subasree, J. Arockia Selvi, M. Arthanareeswari, and Renjith S. Pillai</u> EFFECTS OF PYROLYSIS TEMPERATURE ON THE COMPOSITION OF LIQUID SMOKE DERIVED FROM OIL PALM EMPTY FRUIT BUNCHES - M. Faisal , Asri Gani, Farid Mulana, Hera Desvita and S. Kamaruzzaman BINUCLEAR TRANSITION METAL COMPLEXES DERIVED FROM 3, 3'-DIHYDROXY BENZIDIENE AND 2-AMINO THIO PHENOL: SPECTROSCOPIC, THERMOGRAVIMETRIC, DNA CLEAVAGE, AND ANTIMICROBIAL STUDIES <u>— Kuttiyapillai Sivakumar and Venkatachalam Chandrasekaran</u> FORMULA COMPARISON OF NANOEMULSION AND CREAM CONTAINING MICONAZOLE NITRATE: PENETRATION TEST

FORMULA COMPARISON OF NANOEMULSION AND CREAM CONTAINING MICONAZOLE NITRATE: PENETRATION TEST USING FRANZ DIFFUSION CELLS AND ANTIFUNGAL ACTIVITY TEST ON Tricophyton mentagrophytes, Microsporum canis AND Candida albicans

<u>– H. L. Maha, and Masfria</u>

RAPID DETECTION OF ASHITABA (Angelica keiskei) HERBAL MEDICINE ADULTERATION USING FTIR AND PRINCIPAL COMPONENT ANALYSIS METHOD <u>— Anne Yuliantini , Fenti Salafiah , Aiyi Asnawi</u> NANOGRAM DETERMINATION OF MNII CATALYST IN THE DEGRADATION OF P-BROMOANILINE BY PERIODATE ION — R. D. Kaushik, Jaspal Singh , Malvika Chawla and Kavita Rawat SYNTHESIS, CHARACTERISATION AND CATALYTIC STUDIES OF NANO ZINC OXIDE-ALUMINA FOR DISPLACEMENT REACTION ON CARBONYLDIIMIDAZOLE — <u>S. Sumathi , M. Balaganesh and K.K. Balasubramanian</u> ANTIOXIDANT AND ANTI-MICROBIAL ACTIVITY STUDY OF SYNTHESIZED COPPER, NICKEL AND ZINC METALS SCHIFF BASE DERIVATIVE OF SALICYLALDEHYDE <u>— J. Panda, L. Adhikari , A.Pal , S.S. Rout , S. Pattanaik and P. Pradhan</u> DEVELOPMENT OF TWO-DIMENSIONAL Mg DOPED ZnO NANO HYBRIDS AS ELECTRODE MATERIALS FOR ELECTROCHEMICAL SUPERCAPACITOR APPLICATIONS <u> – Abisheik John Samuel , A. Deepi , G. Srikesh and A. Samson Nesaraj</u> ANTI-DIABETIC PROPERTY OF GREEN SYNTHESIZED ZINC-OXIDE NANOPARTICLES FROM LEAF EXTRACT OF Chrysanthemum indicum PLANT - Neha Silas, and Reena S. Lawrence PHOTOCATALYTIC ACTIVITY OF TiO2/SiO2 PREPARED FROM SILICA CONTAINED IN VOLCANIC ASH FOR AMMONIA REMOVAL <u>— Е.Т. Wahyuni, S. Suherman , D. Setyawati , R. Puspita , and M.Mudasir</u> SYNTHESIS, MOLECULAR DOCKING AND ANTITUBERCULAR ACTIVITY OF NEW BI HETEROCYCLIC COMPOUNDS ON BENZIMIDAZOLE MOIETY <u>— Dhanaja Kotte , Kumaraswamy Gullapelli , Ravichandar Maroju Ramchander Merugu Brahmeshwari Gavaji</u> THE EXFOLIATION PROCESS OF SAWAHLUNTO COAL INTO GRAPHENE THROUGH THE MODIFIED HUMMER METHOD. <u>- V. Purwandari , S. Gea, B. Wirjosentono , A. Haryono , S. Rahayu and Y. A. Hutapea</u> MICROWAVE-ASSISTED SYNTHESIS OF TETRAZOLE BASED BIPHENYLS DERIVATIVES AND THEIR ANTIMICROBIAL **ACTIVITY**

<u> — D. Ashok, Nalaparaju Nagaraju , M. Ram Reddy , Ravinder Dharavath, K. Ramakrishna and M. Sarasija</u>

PREPARATION OF ANTIBACTERIAL IRON-BASED NANOPARTICLES USING Ruellia tuberosa L. ROOT EXTRACTS AS BIOREDUCTOR

<u>— A. Safitri, M. Ramadhan , and A. Sabarudin</u>



OPTIMIZATION SYNTHESIS FATTY ACID ETHYL ESTER AS BIODIESEL FROM PALM FATTY ACID DISTILLATE USING SO42?/ TIO2 CATALYST SUPPORTED BY MESOPOROUS SILICA



<u>— J. Manga, A. Ahmad , P. Taba , and Firdaus</u>

ENHANCEMENT OF ANTIBACTERIAL ACTIVITY OF SUEDE LEATHER THROUGH COATING SILVER NANOPARTICLES. SYNTHESIZED USING PIPER BETLE



<u> E. Rohaeti, E. Kasmudjiastuti , R. S. Murti , and D. Irwanto</u>

LEAF EXTRACT OF Artocarpus altilis [Park.] Fosberg HAS POTENCY AS ANTIINFLAMMATORY, ANTIOXIDANT, AND IMMUNOSUPPRESSANT



<u>– D. H. S. Palupi, D. S. Retnoningrum , M. I. Iwo , and A. A. Soemardji</u>



SYNTHESIS OF (E)-4-METHYL-2-((PHENETHYLIMINO (PHENYL)METHYL)PHENOL AND ITS TRANSITION METAL COMPLEXES, CHARACTERIZATION AND ELECTRICAL CONDUCTIVITY STUDY OF COMPLEXES



– A. B. Sahare and R. B. Mohod

SYNTHESIS OF SOME CHALCONE DERIVATIVES, IN VITRO AND IN SILICO TOXICITY EVALUATION



<u>— A. N. Kristanti , H. Suwito, N.S. Aminah, K.U. Haq, H. D. Hardiyanti, H. Anggraeni, N. Faiza, R.S. Anto and S. Muharromah</u>

QUANTUM MECHANICAL STUDIES ON THE EFFECT OF INTERMOLECULAR ROTATION OF THE STACKING INTERACTION OF **DIAZINE ISOMERS**



— Mahasweta Choudhury, Shruti Sharma, Benzir Ahmed and Bipul Bezbaruah



A NOVEL MICROBIAL FUEL CELL TECHNOLOGY FOR ENERGY GENERATION AND COMPARISON OF POWER DENSITIES FOR DIFFERENT ELECTRODES USING NANOTECHNOLOGY

<u>— Samatha Singh and S. Suresh</u>



SYNTHESIS OF NOVEL PYRAZOLINES AND THEIR ANTIMICROBIAL ACTIVITY



<u>- S. Sathiya , A. Keerthika ,B. S. Krishnamoorthy, S. Nandhabala , S. Aravind , N. Hari and R. Ravikumar</u>



MAROS DISTRICT EXTRACTED USING ULTRASOUND-ASSISTED EXTRACTION



- M. Yasser, M. Rafi , W.T. Wahyuni , S.E. Widiyanti and A.M.I.A Asfar



MOLECULAR DOCKING AND MOLECULAR DYNAMIC SIMULATION OF THE AGLYCONE OF CURCULIGOSIDE A AND ITS

IMPROVEMENT OF CORROSION BY MOLTEN SALTS IN COATING POWDERS MIXED IN DIFFERENT COMPOSITIONS OF

TOTAL PHENOLIC CONTENT AND ANTIOXIDANT ACTIVITIES OF BUNI FRUIT (ANTIDESMA BUNIUS L.) IN MONCONGLOE

DERIVATIVES AS ALPHA GLUCOSIDASE INHIBITORS



<u>— Nursamsiar, M. Siregar , A. Awaluddin , N. Nurnahari , S. Nur , E. Febrina and A. Asnawi</u>



<u> – J. Bautista-Ruiz , A. Chaparro and J.C. Caicedo</u>

YSZ-Al2O3

THERMO-PHYSICAL, SPECTRAL EVALUATION OF MOLECULAR INTERACTIONS IN LIQUID BINARIES OF DIETHYL MALONATE AND AMIDES AT TEMPERATURES (303.15, 308.15, 318.15, 318.15) K



<u>— Ch. Udayalakshmi , Shaik Beebi , P.B. Sandhya Sri ,V.N.S.R. Venkateswararao, G. Lakshmana Rao, and C. Rambabu</u>

SYNTHESIS OF CO-NH2/MESOPOROUS SILICA BIFUNCTIONAL CATALYST USING SIDOARJO MUD AND BOVINE BONE GELATIN TEMPLATE FOR CONVERSION OF USED COOKING OIL INTO BIOFUEL



<u>– W. Trisunaryanti, Triyono and D. A. Fatmawati</u>

DEVELOPMENT OF COMPLEX FERTILIZER TECHNOLOGY WITH IMPROVED AGROCHEMICAL PROPERTIES USING HYDROGEL



<u> – Raikhan ZH. Omirova, Aidar A. Bolysbek , Shavkat D. Shirinov and Abdulakhat T. Dzhalilov</u>



EFFECT OF OKRA (Abelmoschus esculentus L.) FRUIT EXTRACT IN IMPROVING INSULIN SENSITIVITY BY MODIFYING GLUCOSE-REGULATING GENE EXPRESSION



<u>— W. Aligita, S. Muhsinin , K.T. Wijaya , A. Artarini and I.K. Adnyana</u>



Zno incorporated on natural zeolite for photodegradation of methylene blue



<u>— N. P. Diantariani, I. Kartini , A. Kuncaka , and E. T. Wahyuni</u>

PREPARATION OF S-DOPED TIO2 VIA SOL-GEL/ HYDROTHERMAL METHOD AND ITS ACTIVITY AS PHOTODEGRADATION OF RB05



<u> – Adiansyah, I. Elisabeth Purba , M. Tarigan , Lisnadiyanti , Yusnaidar and I.P. Mahendra</u>



EXTRACTION OF Co(II) AND Cu(II) FROM WASTE MATERIALS WITH SEMICARBAZONE OF 4- NITROBENZALDEHYDE



<u>— Jaspal Singh, R. D. Kaushik, Payal Devi and Preeti Das</u>

INFLUENCE OF THE DIFFERENT FORMS OF CRUM RUBBER PARTICLE IN NAO BRAKE PAD MATERIALS



- Adnan Ibrahim Shihab

THE EFFECT OF NICKEL CONTENT IMPREGNATED ON ZEOLITE TOWARD CATALYTIC ACTIVITY AND SELECTIVITY FOR HYDROTREATING OF CASHEW NUT SHELL LIQUID OIL



<u>– M. L. Permata, W. Trisunaryanti , I. I. Falah, M. T. Hapsari and D. A. Fatmawati</u>

Important Links

— About Rasayan

Home
About us
Contact
Archive
Abstracted
Announcement

Editorial Board
Author's Guidelines
Subscription
Manuscript Submission
Current Issue

Rasayan J. Chem. ISSN: 0974-1496(Print) ISSN: 0976-0083(Online) CODEN: RJCABP Frequency: March, June, September & December

© Copyright 2020 RJC. All rights reserved. website designed and managed by ubijournal.



RASAYAN J. Chem.



Vol. 13 | No. 1 | 739 – 746 | January - March | 2020 ISSN: 0974-1496 | e-ISSN: 0976-0083 | CODEN: RJCABP http://www.rasayanjournal.com http://www.rasayanjournal.co.in

EFFECT OF OKRA (Abelmoschus esculentus L.) FRUIT EXTRACT IN IMPROVING INSULIN SENSITIVITY BY MODIFYING GLUCOSE-REGULATING GENE EXPRESSION

W. Aligita^{1,*}, S. Muhsinin¹, K.T. Wijaya¹, A. Artarini² and I.K. Adnyana²

¹ Faculty of Pharmacy, Bhakti Kencana University, Bandung, Indonesia ² School of Pharmacy, Bandung Institute of Technology, Bandung, Indonesia *E-mail: widhya.aligita@bku.ac.id

ABSTRACT

The previous study showed that okra fruit extract improved insulin sensitivity. Insulin resistance played an important role in the pathophysiology of type 2 diabetes. At the molecular level, impaired insulin sensitivity could be caused by several factors, mostly the occurrence of a mutation or posttranlation modification at the insulin receptor or its downstream effector molecule. This study aimed to evaluate the effect of okra fruit extract on glucose regulating gene expressions, such as PPAR-γ, adiponection, resistin, GLUT-4, and GLP-1 gene. Evaluation of this gene expression was performed using the PCR-Gel Base method with the β-actin gene as an internal control gene. The area under the curve (AUC) from the image was determined by using *ImageJ* software. The results showed that the administration of metformin or okra fruit extracts affected the AUC value compared to the positive control, which meant that the expression of each gene changed. Increased expression occurred in PPARG, adiponectin, GLUT-4, and GLP-1 genes; while resistin gene expression decreased. It was concluded that okra fruit extract could improve insulin sensitivity by modifying the expression of PPAR-γ, adiponection, resistin, GLUT-4, and GLP-1 gene.

Keywords: Diabetes Mellitus, *Abelmoschus esculentus* L., Okra Fruit, Insulin Resistance, PPAR- γ

© RASĀYAN. All rights reserved

INTRODUCTION

Diabetes mellitus (DM) is a group of metabolic disorders characterized by hyperglycemia and abnormality of carbohydrate, fat and protein metabolism; caused by the inability of the pancreas to produce insulin or the body's inability to use insulin optimally.^{1,2}

Okra (*Abelmoschus esculentus* (L.) Moench) is belonged to *malvaceae* family. The activity of okra fruit including antioxidant ^{3,4}, lipid inhibitor⁵, antidiabetic ^{6,7,8}, and glucose absorption inhibitor ⁹ as well as a natural absorbent technology in water treatment. ¹⁰

As an antidiabetic, okra fruit extract had a mechanism of action in increasing insulin secretion and sensitivity as well as inhibit glucose absorption in the gastrointestinal tract. The okra fruit extract contained steroids/triterpenoids and flavonoids. Another study showed that the okra fruit rich in α -cellulose and hemicellulose, both belonged to the fiber group. And it was known that fiber could be so helpful to stabilize blood sugar level by suppressing sugar absorption in the intestinal tract. Polyphenols compounds could modulate carbohydrate and lipid metabolism, reduce hyperglycemia, improve adipose tissue metabolism, dyslipidemia and insulin resistance, and stress-sensitive signaling pathways and alleviate oxidative stress and inflammatory processes. While triterpenoids had inhibited the formation of advanced glycation end products and strong antioxidant activity. Antioxidant activity of these compounds is also able to prevent the occurrence of metabolic syndrome.

Decreased insulin sensitivity, also known as insulin resistance syndrome, plays an important role in type 2 diabetes pathophysiology, including its relationship with other symptoms of metabolic syndrome. Insulin resistance is usually associated with the inability of insulin to perform its function in glucose uptake,



metabolism, and storage.¹⁴ At the molecular level, impaired insulin sensitivity can be caused by several factors, such as the occurrence of a mutation or posttranlation modification at the insulin receptor or its downstream effector molecule.¹⁵ Some of the glucose regulating genes include gamma-activated activator receptor peroxisome (PPAR-γ), resistin, GLP-1, GLUT-4, and adiponectin. Peroxisome Proliferator Activated Receptor-gamma (PPAR-γ), adiponectin, and resistin are known to play an important role in glucose transport¹⁶; GLP-1 in stimulating the hormone insulin in the pancreas; whereas GLUT-4 affect glucagon in regulating blood sugar levels.^{17,18}

One of the insulin receptors that started to become the focus in the treatment of diabetes was PPAR- γ . 19,20,21 PPAR- γ , belonged to the PPARs, is a nuclear hormone receptor that acted as a second messenger in regulating glucose uptake and utilization. PPARs consist of PPAR- α , PPAR- δ , and PPAR- γ . The primary effect of PPAR- α is to improve plasma lipid, PPAR- δ responsible for fatty acid oxidation and immunomodulator, while the PPAR- γ improves insulin sensitivity. 19,21 PPAR- γ is highly expressed in fat tissue compared to muscle and liver. However, PPAR- γ activation will increase insulin sensitivity in the muscle and liver by increasing adiponectin and decreasing resistin, leptin, interleukin and TNF- α , and modifying fatty acid uptake and lipolysis, increasing glucose uptake in skeletal muscle and reducing hepatic glucose production and accumulation of liver fat. 15,22

Adiponectin is a peptide hormone, mainly produced by adipocytes, plays a role in glucose regulation and catabolism of fatty acid. Adiponectin levels will increase when the body's fat decrease. Adiponectin level elevation improves insulin sensitivity because the serum adiponectin level has a positive correlation with insulin sensitivity. Resistin (resistant to insulin) is identified as having an association with obesity and insulin resistance. Resistin (resistant to insulin) is identified as having an association with obesity and insulin resistance. The mechanism of resistin in causing insulin resistance is by inhibiting the insulin stimulated-glucose uptake and inhibiting the differentiation of adipose cells. Glucose uptake in muscles and adipose cells are determined by GLUT-4 expression. GLUT-4 is one of the glucose transporter present in muscle and adipose tissue and plays a role in insulin-mediated glucose uptake. GLUT-4 diffuses glucose into muscle and fat cells. GLUT-4 expression in skeletal muscle, heart, and fat cells has decreased in patients with insulin resistance. Another gene that plays a role in glucose regulation is GLP-1. GLP-1 is a polypeptide with a length of 30 amino acids produced in the endocrine part of intestinal epithelial L cells as a product of glucagon gene expression. GLP-1 is a powerful stimulant for insulin and glucagon inhibitors but is quickly converted into inactive metabolites by the DPP-4 enzyme.

The previous study evaluated the okra fruit extract in improving the insulin sensitivity⁸. The study was conducted by administrating okra fruit extract at various doses (25, 50, 100, and 200 mg/kg bw) to fat emulsion induced mice for 14 days. The parameter was the value of the insulin tolerance test constant (C_{ITT}). The result showed a significant insulin sensitivity improvement in the group treated with okra fruit extract at a dose of 50 and 100 mg/kg bw, comparable to metformin as a standard drug. This current study aimed to evaluate the mechanism of action of the okra fruit extract on glucose regulating genes expression responsible in increasing insulin sensitivity.

EXPERIMENTAL

Identification and Authentication of Plant Material

Okra (*Abelmoschus esculentus* L.) fruit was purchased from Research Station for Spices and Medicinal Plants, Bogor, West Java, Indonesia and for its identification and authentication was performed by the Center for Plant Conservation Botanic Gardens, Bogor, West Java, Indonesia.

Preparation of Abelmoschus esculentus L. Extract

After cleaned and cut into pieces, okra fruit was mashed using a blender and the mixture was separated using a filter paper and the filtrate was collected and concentrated using a freeze drier.

Animals

Male Swiss-Webster mice 2-3 months old and weighing 20-30 g were kept at standard laboratory conditions at humidity 70-75%, 24-26°C, and 12 hours light/dark cycle. Animals were fed with water *ad libitum* and standard chow. The methods in this study were carried out by ethics and guide for animal

care and used. Ethical approval was published by the Ethics Committee Padjajaran University (Registration number: 0818020269).

General Procedure

Treatment of Insulin Resistance Animal Model

There were two stages in antidiabetic testing *in vivo*. In the first stage, mice were induced using lipid emulsion⁸ at a dose of 0.42 ml/20 g bw for 2 weeks. In the second stage, mice with insulin resistance (from first stage) were randomly grouped into 6 groups: standard drug group (metformin at dose of 135 mg / kg bw), positive control group, and extract groups at dose of 25 mg/kg bw, 50 mg/kg bw, 100 mg/kg, and 200 mg/kg bw. Administration of metformin and extracts was performed for 14 days. At the end of the study, mice were sacrificed and the adipose tissue and perirenal fat were collected to be evaluated the expression of the PPAR- γ (perirenal fat), adiponectin (adipose tissue), resistin (perirenal fat), GLUT-4 (adipose tissue), and GLP1 gene (adipose tissue).

Semi-Quantitative Gene Expression Evaluation

Primers were derived from the resistin, GLUT-4, PPARG, adiponectin, and GLP1 genes (Table-1).

Gene Name	Gene Code	Table-1: Primers for the Genes Primer	Size (bp)
Gene Ivanie	Gene Code	Timei	Size (op)
Adiponectin ²⁹	Adipoq	Adipoq F: 5'-AGCCT GGAGA AGCCG CTTAT-3'	151
		Adipoq R: 5'-TTGCA GT AGA ACTTG CCAGT GC-3'	
GLP-1R ³⁰	glp1r	Glp-1r F: 5'-TCAGAGACGGTGCAGAAATG -3'	158
		Glp-1r R: 5'- CAGCTGACATTCACGAAGGA-3'	
GLUT-4 ³¹	slc2a4	slc2a4 F : 5'-CTGTGCCATCTTGATGACCGTG-3'	196
		slc2a4 R : 5'-GTTGGAGAAACCAGCGACAGC-3'	
PPAR-γ ³²	pparg	Pparg F : 5'-CCACCAACTTCGGAATCAGCT-3'	434
		Pparg R : 5'- TTTGTGGATCCGGCAGTTAAGA-3'	
Resistin ²⁹	retn	retn F: 5'-TCAAC TCCCT GTTTC CAAAT GC-3'	151
		retn R: 5'-TCTTC ACGAA TGTCC CACGA-3'	

Table-1: Primers for the Genes

Primer quality was evaluated using the BLAST software (Basic Local Alignment Search Tool) through http://blast.ncbi.nlm.nih.gov/Blast.cgi. The alignment step was performed by using software (http://ncbi.nlm.nih.gov/tools/primer-blast/) by entering the nucleotide sequences. The next step was calculating the product length based on the forward and reverse primer used.

The expression of the genes at the mRNA level was carried out using the PCR gel base method with band thickness and area under the curve on the electropherogram as the parameters. Total RNA from each organ was isolated using SV Total RNA Isolation System kit. Cell breakdown was carried out by grinding each organ sample using mortar pestle and liquid nitrogen, then adding 175µl of RNA Lysis solution (particularly for liver used 350 µl solution), 350 µl RNA Dilution Buffer. The mixture was incubated at 70°C for 3 minutes, then centrifuged for 10 minutes at a speed of 13000 rpm. An 95% ethanol was added to the supernatant in the spin column as a precipitation process. Total RNA was washed using 600 µl RNA wash solution and centrifuged for 1 minute at 13000 rpm. The next step was the removal of Genomic DNA by adding 40 µl Yellow core buffer, MnCl 2 5 µl and 5 µl DNAse I enzyme and incubated at a temperature of 20-25°C for 15 minutes. The reaction was stopped by adding 200 µl DNAse Stop Solution to the mixture. Total RNA was re-washed using 600 µl and 250 µl RNA wash solution and centrifuged for 2 minutes at a speed of 14500 rpm at each step. Total RNA was eluted using 100 µl Nuclease Free Water and centrifuged for 1 minute at 13000 rpm. The isolated total RNA could be stored in the elution tubes at a temperature of -70°C (as an alternative could be stored at temperatures of -20 to -30°C).

The procedure of the RT-PCR method from isolated RNA was performed as stated in the AccessQuick TM RT-PCR System (Promega) kit protocol. The incubation temperature of RNA when transcripted into cDNA was 45°C for 45 minutes. The cDNA was then amplified by 35 cycles, including the initial denaturation stage at 95°C for 2 minutes then continued with annealing stage at 57°C for 30 seconds, Extention stage at 72°C for 30 seconds, and final extension at 72°C for 5 minutes. The amplification product was stored at 4°C.

The amplification (amplicon) product was then detected using agarose gel electrophoresis. The electrophoresis process was run at 100 volts for 30 minutes. Electrophoregram was then visualized under planar UV and photographed using a camera with an ISO value 3200; $E \frac{1}{6}$; magnification 37.00 mm; and $E \frac{f}{5.6}$ using digiCam software. The area under the curve (AUC) from the image was determined using ImageJ software. This value interpreted the total gene expression and DNA in intact conditions because it was directly proportional to the band's thickness.

RESULTS AND DISCUSSION

The PCR technique can generally be used with certain modifications for gene analysis and the results can be directly applied. The success of a PCR process depends on the primer. The primers in the PCR process serve as a barrier to the target DNA fragment to be amplified (forward primer) and provide a hydroxy (-OH) group at the 3 'end that is required for DNA extension (reverse primer). The primer design was carried out based on known DNA sequences then evaluated the suitability of the nitrogen base sequence using the GenBank database system on the NCBI website. The primer obtained from the literature were evaluated see the primary quality using the **BLAST** software to http://ncbi.nlm.nih.gov/tools/primer-blast/. BLAST is a feature in NCBI which functions to analyze whether there are primary similarities between the primer used with genes in musculus on GenBank DNA and find out other nucleotide sequences that are similar to their primary sequence. The result showed that the primer sequence used specifically in the mus musculus was by the genes to be evaluated.

The AUC from the electropherogram was shown in Fig.-1. The results of the previous experiment showed that the administration of metformin or okra fruit extract increased insulin sensitivity. In this current experiment, we will evaluate the effect of okra fruit extract in influencing the expression of PPARG, adiponectin, resistin, GLUT-4, and GLP1 genes. These genes play an important role in glucose regulation and determine the level of insulin sensitivity.

From Fig.-1, the positive control group showed reduced PPARG gene expression compared to the negative control group. This means that fat emulsion administration could reduce insulin sensitivity by inhibiting PPARG gene expression. Meanwhile, the group given metformin showed an increase in PPARG gene expression. The same results were also shown in the group given okra fruit extract. PPAR-γ is found in high amounts in fat cells and a small portion in muscle cells such as adipose tissue. The activation of PPAR-γ will stimulate the process of adipogenesis, differentiation and distribution of adipocytes so that it stimulates the efficiency of energy storage and mediates the metabolic effects of insulin and urges anti-inflammatory action. PPARG activation can also increase adiponectin and decrease resistin. Otherwise, in the condition of obesity or insulin resistance, reduction in PPARG expression will decrease adiponectin expression and an increase in resistin. This is consistent with the results of experiments that showed a decrease in adiponectin gene expression and an increase in resistin gene expression.

The adipoq gene that encodes adiponectin is an adipokin group that is usually expressed under certain conditions. Conditions that affect the level of adipoq gene expressions such as type 2 diabetes mellitus accompanied by insulin resistance and obesity. Adiponectin is present in adipose and plasma tissue, but in conditions of obesity, adiponectin can also be detected in the liver, pancreas and other fatty tissues such as perirenal tissue. Adiponectin expression has the opposite effect with other adipokines such as resistin and leptin. Adiponectin in a person with DM type 2 and obese conditions, will be lower than normal conditions (non-obese) or DM type 1. This gene has a role to prevent the occurrence of insulin resistance, especially in type 2 DM by providing the effect of increasing insulin sensitivity. Adiponectin influences the process of gluconeogenesin in the liver, urging the effect of insulin sensitivity on skeletal

muscles and the liver thereby increasing transport and burning of fatty acids. Increased burning of these fatty acids will also increase energy in skeletal muscles and cause a decrease in triglycerides in skeletal muscle and increase insulin transduction.³⁶




Fig.-1: AUC from Electrophoregram

Figure-1 showed the Adipoq gene in the positive control group had lower AUC compared to the negative control group and compared to the metformin and extract group. This means that the administration of fat emulsion caused a decrease in the expression of adiponectin, which in turn caused insulin resistance. Treatment with metformin or okra fruit extracts would increase the expression of the adiponectin gene so that it could improve insulin sensitivity.

Resistin expression is affected by PPARG activation. Resistin and other adipokines, such as leptin and TNF-α, will generally increase in conditions of insulin resistance and obesity. This elevation in resistin expression will be associated with type 2 DM. The result showed that the positive control group experienced type 2 DM characterized by increased RETN gene expression. The difference in resistin expression could be seen in the treated groups. In the metformin and okra fruit extract groups, resistin expression decreased. This indicated that there was an improvement in insulin sensitivity in animal models. The results showed that there was a link between PPARG gene activation in its role in increasing adipoq gene expression and decreasing the expression of resistin genes.

GLUT-4 is an insulin receptor located on the surface of cell membranes, precisely in muscle and fat tissue. When insulin resistance occurs, GLUT-4 loses its stimulation to diffuse glucose into muscle cells, so in this condition, DM sufferers feel weak because there is no energy intake into the muscles. This is also influenced by decreased expression of the adiponectin gene. PPAR-γ expression can stimulate the expression of GLUT-4.³⁷ The results showed that the induction of high-fat emulsion was very influential on GLUT-4 gene expression, as well as treatment with metformin or extract. The positive control group showed lower GLUT-4 gene expression when compared to the negative control group. This means that GLUT4 gene expression reduction had a decreased effect on insulin sensitivity. Whereas in the group

given treatment, both with metformin or okra fruit extract, it could be seen that an increase in GLUT4 gene expression also affected the improvement of insulin sensitivity.

GLP-1 is part of the hormone incretin which has a contribution to insulin release. Endocrine L cells of the intestinal epithelium produce GLP-1 as a product of glucagon gene expression and are stored in the pancreas and present in plasma. The expression of this gene will indirectly indicate the condition of the pancreas. A healthy pancreas will produce enough GLP-1 gene expression to stimulate insulin. Conversely, GLP-1 gene expression will decrease under DM conditions. This is related to the presence of the DPP-4 enzyme whose task is to convert GLP-1 into inactive metabolites, which causes a decrease in insulin release and increased release of glucagon so that glucose in the blood is not transported into cells. The test results showed GLP1R gene expression on positive control was lower than the negative control, and the administration of metformin or okra fruit extracts could increase the expression of these genes. This showed that GLP1 gene expression reduction contributed to insulin resistance, and treatment using metformin or okra fruit extracts would increase GLP1 gene expression and in the end improve insulin sensitivity.

Based on experimental results, it was known that insulin resistance was largely determined by the expression of glucose-regulating genes. Administration of fat emulsion as an induction caused a decrease in insulin sensitivity characterized by a decrease in the insulin tolerance test constant (C_{ITT}).⁸ At the molecular level, reduction in insulin sensitivity was caused by changes in the expression of glucose-regulating genes, in this case, a decrease in PPARG, adiponectin, GLUT4, and GLP1 gene expression, and an increase in the expression of resistin genes. Metformin as the first line of type 2 DM therapy showed the effect of improving glucose metabolism and diabetes-related complications. However, the mechanism of action in detail is not fully understood. Physiologically, metformin can reduce hepatic glucose production.³⁹ Molecularly, metformin can increase the expression of PPARG⁴⁰, adiponectin⁴¹, GLUT-4³⁷, GLP1⁴², and decrease the expression of resistin genes⁴³, as shown in Fig.-1. Similar results were also shown in the group administered okra fruit extracts; although in some genes, the changes given were not exactly like metformin. From Fig.-1, it could also be seen that increasing the dose caused an increase in the effect of affecting the expression of these genes.

CONCLUSION

Okra fruit extract improved insulin sensitivity in insulin resistance animals by increasing the expression of PPAR- γ, adiponectin, GLUT-4, and GLP-1; and reducing the expression of resistin gene.

ACKNOWLEDGMENT

The authors would like to thank The Ministry of Research, Technology, & Higher Education Republic of Indonesia, for fully supporting this research, through the Research of Inter-university Cooperation scheme in 2019.

REFERENCES

- 1. B. G. Wells, J. T. Dipiro, T.L.Schwinghammer, C. V. Dipiro, Pharmacotherapy Handbook. McGraw-Hill Education 2015.
- 2. American Diabetes Association, 38 (Suppl. 1), S8(2015), DOI:10.2337/dc15-S005
- 3. G. Shui and L.L. Peng, *Journal of Chromatography*, **1048**, 17(2004), **DOI:**10.1016/j.chroma.2004.07.032
- 4. S.E. Atawodi, J.C. Atawodi, G.A. Idakwo, B. Pfundstein, R. Haubner, G. Wurtel, Spiegelhalder and R.W. Owen, *Journal of Medicinal Food*, **12(6)**, 1316(2009), **DOI:**10.1089/jmf.2008.0211
- 5. N.M. Ansari, L. Houlihan, B. Hussain, A. Pieroni, *Phytotherapy Research*, **19(10)**, 907(2005), **DOI**:10.1002/ptr.1756
- 6. V. Sabitha, S. Ramachandran, K.R. Naveen, and K. Panneerselvam, *Journal of Pharmacy and Bioallied Sciences.*, **3(3)**, 397(2011), **DOI:**10.4103/0975-7406.84447
- 7. N. Mishra, D. Kumar, S.I. Rizvi, *Journal of Dietary Supplements*, **13(6)**, 540, 634 (2016), **DOI**:10.3109/19390211.2016.1164787
- 8. W. Aligita, S. Muhsinin, E. Susilawati, Dahlia, D.S. Pratiwi, D. Aprilliani, A. Artarini, I.K. Adnyana, *Rasayan Journal of Chemistry*, **12(1)**, 157(2019), **DOI:**10.31788/RJC.2019.1215059

- 9. E. Abi, I. Abi, & M.J. Ladan, *Endocrinology and Diabetes Research Journal*, **3(1)**, (2017), **DOI**:10.4172/2470-7570.1000118
- 10. S. Ramesh, J.S. Sudarsan, M. Jothilingam, Rasayan Journal of Chemistry, 9(3), 325(2016)
- 11. R. Testa, A.R. Bonfigli, S. Genovese, V.D. Nigris, & A. Ceriello, Nutrients, 8(5), 310 (2016), DOI:10.3390/nu8050310
- 12. J. Nazaruk & M. Borzym-Kluczyk, Phytochemistry Reviews, **14(4)**, 675(2015), **DOI:**10.1007/s11101-014-9369-x
- 13. G. Haro, I. Iksen, R.M. Rumanti, N. Marbun, R.P. Sari, R.P.J. Gultom, *Rasayan Journal of Chemistry*, **11(1)**, 232(2018), **DOI:** 10.7324/RJC.2018.1112011
- 14. B.B. Kahn & J.S. Flier, *The Journal of Clinical Investigation*, **106(4)**, 473(2000), **DOI:**10.1172/JCI10842
- 15. O. Pedersen, J.F. Bak, P.H. Andersen, S. Lund, D.E. Moller, J.S. Flier, B.B. Kahn, *Diabetes*, **39**, 865(1990), **DOI:**10.2337/diab.39.7.865
- 16. P.M. Kumar, M.V. Venkataranganna, K. Manjunath, G.L. Viswanatha and G. Ashok, *Journal of Intercultural Ethnopharmacology*, **5(2)**, 146(2016), **DOI:**10.5455/jice.20160224051727.
- 17. A.E. Civitarese, C.P. Jenkinson, D. Richardson, M. Bajaj, K. Cusi, S. Kashyap, R. Berria, R. Belfort, R.A. DeFronzo, L.J. Mandarino and E. Ravussin, *Diabetologia*, **47(5)**, 816(2004), **DOI**:10.1007/s00125-004-1359-x
- 18. D. Le Roith & Y. Zick, *Diabetes Care*, **24(3)**, 588(2001), **DOI:**10.2337/diacare.24.3.588
- 19. M.B. Wright, M. Bortolini, M. Tadayyon, & M. Bopst, *Molecular Endocrinology*, **28(11)**, 1756(2014), **DOI:**10.1210/me.2013-1427
- 20. R. Rios-Vazquez, R. Marzoa-Rivas, I. Gil-Ortega and J.C. Kaski, *American Journal of Cardiovascular Drugs*, **6(4)**, 231(2006), **DOI:**10.2165/00129784-200606040-00003
- 21. J.P. Wilding, *Diabetes, Obesity, and Metabolism*, **14(11)**, 973(2012), **DOI:**10.1111/j.1463-1326.2012.01601.x
- 22. U. Meier, Clinical Chemistry, 50, 1511(2004), DOI:10.1373/clinchem.2004.032482
- 23. G. Litwack: Adiponectin Volume 90 of Vitamins and Hormone. Academic Press 2012.
- 24. J. Chen, L. Wang, Y.S. Boeg, B. Xia and J.J. Wang, *Endocrinology*, **175(2)**, 499(2002), **DOI**:10.1677/joe.0.1750499
- 25. C. McTernan, P. McTernan, A. Harte, P. Levick, A. Barnett and S. Kumar, *The Lancet*, **359**, 46(2002), **DOI:**10.1016/S0140-6736(02)07281-1
- 26. E.E. Kershaw & J.S. Flier, *Journal of Clinical Endocrinology and Metabolism*, **89(6)**, 2548(2000), **DOI:**10.1210/jc.2004-0395
- 27. Martini and Frederic: Fundamentals of Anatomy and Physiology 90th ed. Pearson Education Inc 2012.
- 28. P.R. Sheperd and B.B. Kahn, *The New England Journal of Medicine*, **341(4)**, 248(1999), **DOI**:10.1056/NEJM199907223410406
- 29. S.K. Park, S.Y. Oh, M.Y. Lee, S. Yoon, K.S. Kim and J.W. Kim, *Diabetes*, **53(11)**, 2757(2004), **DOI**:10.2337/diabetes.53.11.2757
- 30. L.Bai, G. Meredith and B.E. Tuch, *The Journal of Endocrinology*, **186(2)**, 343(2005), **DOI:**10.1677/joe.1.06078
- 31. D. Furuya, A.C. Poletto, H.S. Freitas and U.F. Machado, *Journal of Molecular Endocrinology*, **49(2)**, 97(2012), **DOI:**10.1530/JME-12-0037
- 32. M.Inoue, T. Ohtake, W. Motomura, N. Takahashi, Y. Hosoki, S. Miyoshi, Y. Suzuki, H. Saito, Y. Kohgo and T. Okumura, *Biochemical Biophysical Research Communications*, **336(1)**, 215(2005), **DOI**: 10.1016/j.bbrc.2005.08.070
- 33. D.A. Hewajuli and N.L. Dharmayanti, *Indonesian Bulletin of Animal and Veterinary Sciences*, **24(1)**, 16(2014), **DOI:**10.14334/wartazoa.v24i1.1022.
- 34. M.Boyraz, F. Cekmez, A. Karaoglu, P. Cinaz, M. Durak, A. Bideci, *Biomarkers in Medicine*, **7(5)**, 737(2013), **DOI:**10.2217/bmm.13.13
- 35. A.A. Gumbs, Obesitiv Surgery, 18(9), 1183(2008), DOI:10.1007/s11695-008-9599-3

- 36. J. Chen, H. Zhao, X. Ma, Y. Zhang, S. Lu, Y. Wang, C Zong, D. Qin, Y. Wang, Y. Yingfeng Yang, X. Wang and Y. Liu, *Cellular Physiology and Biochemistry*, **42(3)**, 1165(2017), **DOI:** 10.1159/000478872
- 37. Z. Wu, Y. Xie, R.F. Morrison, N.L. Bucher, S.R. Farmer, *Journal of Clinical Investigation*, **101(1)**, 22(1998), **DOI:**10.1172/JCI1244
- 38. M.A. Nauck, I. Vardarli, C.F. Deacon, J.J. Holst, J.J. Meier, *Diabetologia*, **54(1)**, 10(2011), **DOI:** 10.1007/s00125-010-1896-4
- 39. G. Rena, D.G. Hardie, E.R.P. Pearson, *Diabetologia*, **60(9)**, 1577(2017), **DOI:**10.1007/s00125-017-4342-z
- 40. E.M. Elia, C. Pustovrh, S. Amalfi, L. Devoto, A.B. Motta, *Fertility and Sterility*, **95(8)**, 2534(2011), **DOI**:10.1016/j.fertnstert.2011.02.004
- 41. G. Basios, E. Trakakis, Ch. Chrelias, P. Panagopoulos, V. Vaggopoulos, P. Skarpas, D. Kassanos, G. Trimidiadis, E. Hatziagelaki, *Gynecological Endocrinology*, **31(2)**, 136(2015), **DOI:** 10.3109/09513590.2014.975684
- 42. E. Bahne, E.W.L. Sun, R.L. Young, M. Hansen, D.P. Sonne, J.S. Hansen, U. Rohde, A.P. Liou, M.L. Jackson, D. de Fontgallant, P. Rabbit, P. Hollington, L. Sposato, S. Due, D.A. Wattchow, J.F. Rehfeld, J.J. Holst, D.J. Keating, T. Vilsboll, F.K. Knop, *Journal of Clinical Investigation (JCI) Insight*, 3(23), (2018), DOI:10.1172/jci.insight.93936
- 43. H.Fujita, H. Fujishima, T. Morii, J. Joshimura, M. Kakei, S. Ito, *Biochemical and Biophysiological Research and Communications*, **298(3)**, 345(200), **DOI:**10.1016/s0006-291x(02)02464-6

[RJC-5555/2019]