

Volume 11, Issue 05 May 2019.

- Applications of 3D Printing in Dentistry A Review Pages: 1670-1675 Abarna Jawahar, G.Maragathavalli Abstract | DPF (854 K)
- Assess the Cultural Practices on Newborn Care among Mothers Residing in Kayamkulam Pages: 1676-1678
 Dr. Kala Barathi S, Ms.Surumi.M
 Abstract | DPF (754 K)
- 3. Analysis of changes in market characteristics of Essential Medicines within the frames of state program of increasing availability of medicines in Ukraine Pages: 1679-1686 Oleh Samborskyi, Mykola Slobodyanyuk, Hanna Panfilova, Viacheslav Lebedynets, Natalia Bogdan Abstract | DPF (1024 K)
- Chronopharmacological dependence of the anticytolytic and antioxidant influence of silymarine and arginine glutamate under conditions of drug-induced hepatitis in rats
 Pages: 1687-1692
 O.Ya. Mishchenko, K. O. Kalko, S. M. Drogovoz, N. S. Bogdan, O. V. Gerush, L. B. Ivantsyk, A. D. Gordienko

Abstract | Z PDF (879 K)

5. Effect of Drying and Roasting to Antioxidant Property and Stability of Dried Roasted Walnut (Juglans Regia)

Pages: 1693-1697 N. P. Minh, P. T. L. Pha, N. T. T. Trang, B. N. Y <u>Abstract</u> | DF (916 K)

- 6. Production of Pickled Chinese Olive (Canarium Album) Fruit Pages: 1698-1702
 N. P. Minh, P. X. Mai, K. S. Du, T. M. Thang, V. H. Dao, H. P. Quyen Abstract | PDF (854 K)
- 7. Some Variables Affecting To Gooseberry (Phyllanthus acidus) Wine Fermentation *Pages:1703-1709*N. P. Minh, H. Q. Khai, N. T. Loan, L. V. Duc
 Abstract | DPF (1024 K)
- 8. Mixture of Extract Nerium indicum Mill and Tithonia diversifolia (Hemsley) A. Gray. Inhibits the Migration Activity, Expressions of TGF-β1 and VEGF Keloid Fibroblasts *Pages: 1710-1714* Indwiani Astuti, Mae Sri Hartati Wahyuningsih <u>Abstract</u> | DF (916 K)

- 9. Polymorphism of BCL-2 (-938C > A) and associated with breast cancer susceptibility of Iraq population *Pages: 1715-1720* Najwa Shihab Ahmed, Mahmood, Ahmed Muhi, Najlaa, Qassim Muftin
 <u>Abstract</u> | [™] PDF (1100 K)
- Phase and Microelement Composition of the Kidney Stones of Ob' River Watershed Residents Pages: 1721-1726
 Alphing B. Taugaphana, Ilug V. Korolkov, Alayander I. Cubanay, Eugany Vy, Eilatov, Eugany V, Da

Alphiya R. Tsygankova, Ilya V. Korolkov, Alexander I. Gubanov, Evgeny Yu. Filatov, Eugene V. Pechkovskii, Gennadiy Yu. Yaryn and Denis V. Safonov

<u>Abstract</u> | 🔁 <u>PDF</u> (937 K)

11. Development and Validation of UV Spectrophotometric method for in-vitro studies of Anastrozole Invasomes

Pages: 1727-1730 Vidya. K, Lakshmi. PK, Padmavathi. Y Abstract | DF (852 K)

12. Assess the Attitude Regarding Mental Illness among the Community Residents in Meppur

Pages: 1731-1732 Angelin Lavanya. S, Abstract | DF (774 K)

13. Organizational and Economic Justification of Medicamental Provision of the Injured Soldiers with Thoracoabdominal Trauma

Pages: 1733-1741 Andriy Solomennyi, Oleksandr Dobrovolnyi, Natalia Takhtaulova, Mariia Bilous <u>Abstract</u> | DF (1024 K)

14. Study of mutagenicity, embryotoxic action, immunotoxicity and allergenicity of Xenon difluoride *Pages: 1742-1748*

Kanat Sarsenbayev, Mariana Sarsembayeva, Kulzada Lakhanova, Gulnar Kylyshbaeva, Assiya Mamytova, Zharas Berdenov

<u>Abstract</u> | ¹ <u>PDF</u> (1100 K)

15. Research and testing of some grain varieties from Croatia under the agroecological conditions of Kosovo Pages: 1749-1752

Bakir Kelmendi, Vlora Gashi, Ismajl Cacaj, Defrime Berisha, Arsim Elshani, Nazmi Hasnaj Abstract | DF (820 K)

16. **Brusatol- as potent chemotherapeutic regimen and its role on reversing EMT transition** *Pages:1753-1762* Aarthi.R,

Abstract | Z PDF (1400 K)

17. A Brief Outlook on Cancers and Preventive Methods

Pages: 1763-1765 Shilpa Gopal and Thirunavukkarasu Sivaraman Abstract | DF (833 K) 18. Simple Titrimetric, Spectrophotometric and Gravimetric Methods for the Assay of Pitavastatin Calcium in a Green Manner

Pages: 1766-1774 Niranjani S, Venkatachalam K <u>Abstract</u> | DF (1024 K)

 19. Clinical Evaluation of Vaccines Pages: 1775-1780 Janani.M, Nagasamy Venkatesh. D <u>Abstract</u> | DF (826 K)

20. Validated RP-HPLC Method for the Determination of Clofarabine in Bulk and Tablet Dosage Form Pages: 1781-1786

Dr.P.D.Chaithanya Sudha Abstract | 🔁 PDF (1024 K)

21. Comparision of Commercially available Desensitizing Toothpastes in the Management of Dentin Hypersensitivity-A Randomized Controlled Clinical Trial

Pages: 1787-1789 Dr.Devi Bala.R.T, Dr. Dhayanand John Victor, Dr. Sangeetha Subramaniam , Dr. PSG Prakash , Dr. Abirami.T Abstract | DPF (773 K)

22. Phytochemical Screening, Anti-oxidant and Anti-microbial Activity of leaves of Cleome rutidosperma DC. (Cleomaceae)

Pages: 1790-1795 Pranabesh Ghosh, Maitrayee Biswas, Swagata Biswas, Alolika Dutta, Labani Hazra, Sudip Kumar Nag, Sayantan Sil, Sirshendu Chatterjee

<u>Abstract</u> | 🔁 <u>PDF</u> (868 K)

23. Soybean Oil-Hpmck15M Based Oleohydrogel Hybrid: Novel Approach to Improve Drug Release Pages: 1796-1802

Dipanjana Ash, Sutapa Biswas Majee and Gopa Roy Biswas

<u>Abstract</u> | 🔁 <u>PDF</u> (1024 K)

24. Evaluation of behavioural and cognitive effects against diabetic model in young and adult rats *Pages:1803-1807* Ratna.B, Johnson.C.John

<u>Abstract</u> | 🔁 <u>PDF</u> (797 K)

- 25. Formulation and evaluation of herbal chocolate as nervine tonic *Pages: 1808-1813*Pallavi D. Pawar, Akshada A. Bakliwal, Swati G. Talele, Dr. Anil G. Jadhav Abstract | PDF (1024 K)
- 26. Stability Indicating HPLC Method for Determination of Gatifloxacin in Ophthalmic Dosage Form Pages: 1814-1820

A.Caroline Grace, T.Prabha, M.Jagadeeswaran, K.Srinivasan, T.Sivakumar <u>Abstract</u> | 2 PDF (990 K)

- 27. Antioxidant Activity, Total Phenolic and Flavonoid Contents of Notonia Grandiflora Wall. Extracts Pages: 1821-1825
 Meera Paul, Dr. R. Sanilkumar, Dr. Sabu M.C
 <u>Abstract</u> | DF (926 K)
- 28. Bioactive compounds and antimicrobial activity of fungal crude extract from medicinal plants *Pages:1826-1833* H.Sheeba, M. Syed Ali, V. Anuradha

Abstract | ¹/₂ PDF (1500 K)

29. Touchdown PCR combined with semi degenerate primers for rapid amplification of HOXD9 loci in humans

Pages: 1834-1837 Praveen Kumar Vemuri, Nandini Kollipara, Shamanta Nasika, Ananya Kanneganti, Vijaya Lakshmi Bodiga, Suryanarayana Veeravalli

<u>Abstract</u> | 🔁 <u>PDF</u> (870 K)

30. **Preparation of an Amino Acid Alliin from the Iraqi Garlic and using it as a Substrate for Alliinase** *Pages:1838-1841*

Dhia Falih Al-Fekaiki, Iman Hadi Al-Taai Abstract | DF (860 K)

31. Association of Irisin concentration with clinical Parameters in Obesity patient compared with sport individual in Iraqi population

Pages: 1842-1848 Rahman S. Zabibah, Hanaa Addai Ali, Salwa Jaber Al-Awadi Abstract | DPF (1024 K)

32. An epidemiological, diagnostic and therapeutic study of Giardia lamblia in Anbar province - Iraq Pages: 1849-1855

Suad Shallal Shahatha Abstract | 🔁 PDF (1200 K)

33. Mandibular Degree II Furcation Defect Treatment With 1%Alendronate Gel Alone or In Combination with Platelet- Rich Fibrin

Pages: 1856-1860 Mayssam A.Tahir AL -khaiyat, Saif S.Saliem, Hyder R.Abdulbaqi, Mohened M.Thumad Abstract | DDF (845 K)

34. A Stability Indicating HPTLC Method for Apremilast and Identification of degradation products using MS/MS.

Pages: 1861-1869 Ritesh P. Bhole, Sachin R. Naksakhare, Chandrakant G. Bonde <u>Abstract</u> | DF (1300 K)

35. Knowledge, Attitude & Practice assessment regarding Oral Care in Palliative patients among Dental Graduates

Pages:1870-1873 Shilpa Syam, Prof.Dr.T.N.Uma Maheswari Abstract | DF (887 K) 36. In vitro Antioxidant Activity, Total Phenolic and Total Flavonoid Contents of Dracaena terniflora Roxb. Root Extracts.

Pages: 1874-1879 Kavitha Vasudevan, Dr. R .Sanil Kumar , Dr. M.C Sabu Abstract | DF (971 K)

37. Research Of Rheological Properties Of Ointment With Water-Soluble Protein-Polysaccharide Complex Of Oyster Mushroom

Pages: 1880-1883 O.A. KOLPAKOVA, N.V. KUCHERENKO, H.P. KUKHTENKO Abstract | DF (1100 K)

38. Peri Implant Microflora in Health and Disease Pages:1884-1890
Dr. Sathya Ramanathan, Dr. PSG Prakash, Dr. Devapriya Appukuttan,Dr.Sangeetha Subramanian, Dr.D.J Victor

<u>Abstract</u> | ¹ <u>PDF</u> (903 K)

 Study of Cytotoxic Activity of Two Species of Portulaca on Cancer Cell Lines Pages: 1891-1896
 Trupti P. Durgawale, Dr. Chitra C. Khanwelkar, Dr. Kailas D. Datkhile, Pratik P. Durgawale, Dr. Satish

Abstract | 🔁 PDF (1200 K)

V.Kakade

40. Antihyperlipidemic Activity and HMG CoA Reductase Inhibition of Ethanolic Extract of Zingiber Cassumunar Roxb in Fructose-Induced Hyperlipidemic Wistar Rats Pages:1897-1901

P.Hasimun1, A. Sulaeman, Y. Mulyani, W. Nur Islami, F. Apriany Timika Lubis Abstract | DPF (840 K)

- 41. Plant Derived Vaccines New Door to Periodontal Vaccine Pages: 1902-1906
 Dr.Devi Bala.R.T, Dr. Anupama Tadepalli
 Abstract | DF (902 K)
- 42. Chronopharmacological dependence of the anticytolytic and antioxidant influence of silymarine and arginine glutamate under conditions of drug-induced hepatitis in rats *Pages:1907-1913*O.Ya. Mishchenko, K. O. Kalko, S. M. Drogovoz, N. S. Bogdan, O. V. Gerush, L. B. Ivantsyk, A. D. Gordienko

Abstract | The PDF (877 K)

43. Artifacts in Cone Beam Computed Tomography – A Retrospective study Pages: 1914-1917
Shilpa Syam, Prof. Dr. T.N. Uma Maheswari
<u>Abstract</u> | DF (862 K)

- 44. A Facile Synthetic Method for the Synthesis of new 7',9'-dihydrospiro[indoline-2,11'-pyrazolo[3,4-f]pyrimido[4,5-b]quinoline]-3,8',10'(1'H,6'H)-trione and its BiologicalEvaluation Pages: 1410-1415
 B.Srinivas, J. Suryachandram, P. Sadanandam, M. V. Basaveswara Rao and K. Prabhakara Rao Abstract | DF (1100 K)
- 45. Research of maize hybrids from Croatia of the agro ecological conditions of Kosovo Pages: 1926-1929
 Kelmendi. B, Cacaj I, Shala N, Berisha D
 Abstract | DPF (837 K)
- 46. Acoustic Properties of Binary Liquid Mixtures Containing Dimethyl Formamide with Anilines At Different Temperatures

Pages: 1930-1936 K.Govinda Rao, Shaik.Babu Abstract | 🔁 PDF (1024 K)

 47. A review on biosynthesis, characterization and Antimicrobial effect of silver nanoparticles of Moringa olifera (MO-AgNPs) Pages: 1937-1943 Saurabh Dixit, Astha Tripathi

Abstract | The PDF (913 K)

48. Movement Disorders and Nonmotor Symptoms in Patient with Idiopathic Stiff Person Syndrome Pages: 1944-1947

A.S. Maksimova, O.E. Zinovyeva, O.A. Solokha, A.L. Pilyuzina, D. S Petelin, B. A. Volel, A.D. Ulanova, I.M. Vikhlyantsev

<u>Abstract</u> | 🔁 <u>PDF</u> (870 K)

- 49. Comparing number and intensity of doublecortin-immunolabelled cells in the hippocampus and cerebellum in the postnatal mice brains, in different age groups
 Pages:1948-1953 Zainab Zahid Saadoon
 Abstract | [™] PDF (1400 K)
- 50. Phytochemical and GC-MS analysis of ethyl acetate extract of Ailanthus triphysa leaves Pages: 1954-1959
 Abhishek Kumar Yadav, Zebha Mohamed Ali, Abhishek Biswal R, Vivek pazhamalai
 <u>Abstract</u> | DF (885 K)
- 51. Phytochemical, FTIR and NMR analysis of crude extract of Acacia planifrons seeds Pages:1960-1962 Luvincia Fernando M, Abhishek Biswal R, Vivek Pazhamalai Abstract | DPF (844 K)
- 52. **Pharmacovigilance-A Master Key for Drug Safety** *Pages:1963-1970* Jaani.M , Nagasamy Venkatesh D

<u>Abstract</u> | ¹ <u>PDF</u> (1200 K)

- 53. Augmentation of Attached Gingiva with a Variant of Freegingival Autograft-A Case Report Pages: 1971-1973
 Santhosh Venkatassalapathy Sathya Ramanathan
 <u>Abstract</u> | Abstract | Report (878 K)
- 54. Synthesis, characterization and anticancer activity of some new Tetrazoles derived from Quinazolin-4one

Pages: 1974-1978 Rajaram Arulmozhi, Natarajan Abirami, Helen P. Kavitha, Subramanian Arulmurugan <u>Abstract</u> | DF (1100 K)

55. Study of the Fibrinolytic Parameters in the Different Types of Polycythemia in Al –Nasirya City Pages: 1979-1983

Ahmed Habeeb Radhi , Hiba Jasim Hammood , Iqbal Hanash Dhefer Abstract | The PDF (842 K)

56. Advancement In Chronotherapeutic Drug Delivery System: Marketed Technologies And Current Scenario

Pages: 1984-1989 Kulwinder kaur, Nimrata Seth, Naresh Singh Gill Abstract | DDF (822 K)

57. Article Removed Pages: 1990-1998 xxx, xxx

<u>Abstract</u> | ¹ <u>PDF</u> (1200 K)

58. Predisposing Risk Factors of Acute Coronary Syndrome (ACS): A Mini Review Pages:1999-2002 Varghese TP, Kumar AV

Abstract | DPF (806 K)

59. **Review on Philodendron Species-Plant Seeking For Validation of Its Therapeutic Approaches** *Pages:2003-2006* Pankaj Nainwal

Abstract | Z PDF (808 K)

60. Study of Effect of Solvents on Absorption Characteristics of Rifaximin in Visible Region and Its Estimation in Bulk and Dosage Forms

Pages:2007-2011 P.Ravi Kumar, T Jhansi Chary, K. Sahithi, Amani Baquer Shareef, N Raghavendra Babu Abstract | DF (1100 K)

61. An over view of contract research organization and analysis on aro-cro model Pages: 2012-2015
 Shika Susan Mathew, Dr.G.N. K Ganesh, Phatru Patel
 <u>Abstract</u> | DF (790 K)

- 62. Molecular docking studies of bioactive compounds from Stevia rebaudiana for its anti-cancer activity Pages: 2016-2018
 Maheswara Reddy Mallu, Sandeep Vemula and Rajesh Kumar Kante
 <u>Abstract</u> | DF (879 K)
- 63. The activity of adenosine on some oral bacteria: An in vitro study Pages:2019-2021
 Faehaa Azher Al-Mashhadane, Suhad M.Hamdon, Ghadeer H. Aljader
 <u>Abstract</u> <u>PDF</u> (815 K)
- 64. Assess the effectiveness of buerger allen exercise to reduce pedal edema among the adults with varicose veins at Saveetha medical college and hospital

Pages: 2022-2023 Parimala. L, Vishnu Priya. U Abstract | DPF (767 K)

 65. Optimization and Validation of Resveratrol Using Analytical UV Method Development Pages: 2024-2027 Nita B. Deore, Akshada A. Bakliwal

Abstract | 🔁 PDF (850 K)

66. Effect of Sodium Alginate in Combination with Natural and Synthetic Polymers on the Release of Verapamil HCL from its Floating Microspheres

Pages: 2028-2035 K Saravanakumar, Ashok Thulluru, Ramu Samineni, M Ishwarya, Pommala Nagaveni and Nawaz Mahammed Abstract | DPF (1300 K)

67. Does Long Standing Multi-Nodular Goiter Increase the Risk of Thyroid Cancer? Pages: 2036-2038 Aqeel Shakir Mahmood

Abstract | 🔁 PDF (781 K)

68. Allergic asthma is inversely associated with Helicobacter pylori seropositivity

Pages:2039-2041 Hasanain Abdulhameed Odhar, Mohammed Abbas Hasan, Ali Mahmood Rayshan, Murtada Falah Hasan, Suqayna Habeeb Mahmmod, Huda Kahtan Majid

<u>Abstract</u> | 🔁 <u>PDF</u> (780 K)

69. Relationship between hormonal change and ovarian cyst in buffalo

Pages:2042-2047 Kadhem Mohammed sabaa AL-Jabri and Ahmed Aiad kadhem Al-Jubouri ,Ahmed Kudhair AL-Hamary Haider Mahmooed Jasim

Abstract | 🔁 PDF (980 K)

 70. Effect of Polycystic Ovarian Syndrome on the Lipid Profile and sexual Hormones Pages: 2048-2050
 Eqbal A. Gatea, Israa Abd Alhassan Hamdan, Frhan Ala Allah Eabaid

Abstract | ¹ PDF (753 K)

- 71. In vitro antioxidant activity of new Schiff base ligand and its metal ion complexes Pages:2051-2061
 Hayder Muneam A.M., Mohammed H.A. Al-Amery
 <u>Abstract</u> | DF (2200 K)
- 72. Synthesis and Characterization new metal complexes of heterocyclic units and study antibacterial and antifungal

Pages: 2062-2073 Ali T.Bader, Basim I. Al-Abdaly and Ibtisam K. Jassim Abstract | DF (2800 K)

73. Gene expression and single nucleotide polymorphism of interleukin-6 in asthenozoospermic men: A casecontrol study in Iraqi patients

Pages:2074-2079 Farah Thamer Abdullah, Abdul Kareem A.Alkazaz, Bushra J. Al-Musawi, Saad S. Al-Dujaily <u>Abstract</u> | DF (847 K)

74. Endoparasitic nematohelminthes collected from inside frog Rana ridibunda (Pallas, 1771) in two rivers near Baghdad/Iraq

Pages:2080-2083 Saad Muhi Arif Abstract | DPF (869 K)

75. **In Silico Identification of Drug-Like Inhibitors against Mtb-DHDPS: A Shape-Based Approach** *Pages: 2084-2089* Ajijur Rehman, Salman Akhtar, Mohd Haris Siddigui, Mohd Kalim Ahmad Khan

Abstract | The PDF (1200 K)

76. In-vitro Chick Chorioallantoic Membrane Study of Chitosan Capped 5-Fluorouracil Conjugated Gold Nanoparticles

Pages:2090-2094 Akhila Rajan, P.K. Praseetha, V.N. Ariharan and S. T. Gopu Kumar <u>Abstract</u> | DF (1200 K)

77. A randomized control trial for comparison among Nalbuphine, Tramadol and placebo for treating post anesthetic shivering undergoing spinal anesthesia in Cessarian section *Pages: 2095-2098* Parveez Taneja, Yogesh Bansal, Nidhi Sharma, Kewal Krishan Banotra

Abstract | The PDF (767 K)

- 78. Prospects of Use in Medical Practice of Solidago caucasica Kem.-Nath. *Pages:2099-2101*V.V. Fedotova, D. A. Konovalov
 <u>Abstract</u> | DF (869 K)
- 79. A Study to Identify the Oral Mucosal Deterioration among Adult Tobacco Chewers at Thiruninravur Pages:2102-2104

L.Parimala, N.R. Jenisha Abstract | DF (808 K)

- 80. Use of Herbs among Pregnant Women in Saudi Arabia: A Cross Sectional Study. *Pages:2105-2108* Mohammad F. Zaitoun, Marwah E. Al-Nowis , Faten Y. Alhumaidhan , Hany E. Khalil
 <u>Abstract</u> | DF (775 K)
- 81. Medicine Errors in Dosage Forms and strategies to enhance the auxiliary labeling for Patient Care in Kongu Nadu region Tamil Nadu

Pages:2109-2112 MuthuKumar Andhuvanjseral, Prakash .G, Gayathri .R, SundaraGanapathy .R, Mohan .S Abstract | DF (743 K)



Antihyperlipidemic Activity and HMG CoA Reductase Inhibition of Ethanolic Extract of *Zingiber Cassumunar* Roxb in Fructose-Induced Hyperlipidemic Wistar Rats

P. Hasimun¹, A. Sulaeman¹, Y. Mulyani¹, W. Nur Islami¹, F. Apriany Timika Lubis¹

Pharmacology research group, Bandung School of Pharmacy, Jl. Soekarno Hatta No.754, Bandung, West Java, Indonesia 40617

Abstract

HMG-CoA (3-hydroxyl-3-methyl-glutaryl-coenzyme A) reductase is a liver enzyme that plays an important role in the process of cholesterol synthesis in the liver. Statins are one of the major drug classes for lowering cholesterol through inhibition of HMG-CoA reductase enzyme activity. This study aims to determine the effect of *Zingiber cassumunar* Roxb ethanol extract as anti-hyperlipidemia and its effect on HMG-CoA reductase enzyme in the liver. A total of 30 white Wistar rats were grouped randomly into 6 groups: group 1 (receiving drug carrier), group 2 receiving drug carrier), group 3 (receiving simvastatin 0.9 mg / kg), group 4-5 (receiving extract dose 30, 50, and 100 mg / kg). All groups (except group 2) received 25% fructose in drinking water for 21 days. Day 22 was taken by blood for examination of total cholesterol, HDL, LDL, and triglyceride levels. Then the rats were sacrificed for liver organ isolation and made homogenate liver 20% for HMG CoA reductase enzyme activity test. The results showed that ethanol extract of *Zingiber cassumunar* Roxb dose 30, 50, and 100 mg / kg can decrease total cholesterol, triglyceride, LDL and increase HDL level compared to control group. Extracts of doses of 30, 50, and 100 mg / kg can inhibit the activity of HMG CoA reductase enzyme with a percentage of 11, 20, and 47% inhibition respectively. The results can be concluded that *Zingiber cassumunar* Roxb has anti-hyperlipidemia activity. One of the mechanisms of action is by inhibiting the activity of HMG CoA reductase enzyme.

Keywords: Fructose, HMG-CoA Reduktase, Hyperlipidemia, Simvastatin, Zingiber cassumunar Roxb

INTRODUCTION

Hyperlipidaemia is a condition in which total cholesterol increases, Low Density Lipoprotein (LDL) are high while High Density Lipoprotein (HDL) decreases, or a combination of these disorders [1]. Hyperlipidaemia is a major cause of coronary heart disease. An estimated 17.5 million people die from cardiovascular disease in 2012, with a percentage of 31% of all deaths in the world. The death, estimated at 7.4 million was due to coronary heart disease and 6.7 million was due to stroke [2]. In Indonesia the prevalence of coronary heart disease, heart failure, and stroke is seen to increase with the increasing age of respondents. The prevalence of stroke is similar in males and females [3].

Fructose is a sweetener widely used in various food and beverage products. The excessive use of fructose is reported to cause various health problems including hyperlipidemia. It has been proven by animal studies that 25% of fructose in drinks for 21 days significantly increased hyperlipidemia [3]. Mice receiving food containing 66% of fructose for 15 days experienced hypertension [4]. Fructose can also form acetyl-CoA used in fatty acid synthesis. High-fructose diets are reported to increase the risk of metabolic syndrome [5]. Metabolic syndrome is characterized by hyperglycemia, obesity, dyslipidemia (hypertriglyceride, decreased HDL, elevated cholesterol) and hypertension [6].

Drugs used to manage hyperlipidaemia are classified into five groups: Statin, Fibrat Acid, Resin, Nicotinic Acid (niacin) and Ezetimibe. Simvastatin is a statin agent that works inhibiting HMG-CoA reductase competitively on the process of cholesterol synthesis in the liver. This enzyme plays a role in converting acetyl CoA to mevalonate acid. But the side effects of using simvastatin as a blood cholesterol-lowering drug can lead to toxic myopathy [7]. Therefore, many hyperlipidemia patients prefer to use medicinal plants as an alternative therapy. This is an opportunity for research of natural materials that are considered more safe and efficacious in controlling and lowering cholesterol levels in the blood.

Zingiber Cassumunar Roxb is a plant that has long been used as a traditional medicine to cope with various diseases including hyperlipidemia. This rhizome, empirically used by the people in Indonesia to overcome fever, headache, cough sputum, colds, constipation, jaundice, worm infection, muscle aches and obesity [8].

Based on the above description, this study aimed to determine the antihiperlipidemia activity of ethanol extract from *Zingiber Cassumunar* Roxb rhizome and its effect on HMG CoA reductase enzyme activity that plays an important role in the synthesis of cholesterol in the liver.

MATERIALS AND METHODS

Plant Material

Rhizome of *Zingiber cassumunar* obtained from Manoko Lembang plantation, Bandung, West Java, Indonesia. Furthermore, botanical plant identification was done at the Laboratory of Plant Taxonomy Department of Biology, Padjadjaran University, Bandung (no. 121 / HB / 01/2016).

Extract Preparation

The fresh rhizomes are washed to be cleaned from impurities, then cut into small pieces and dried using 37°C oven. Dry rhizome is made powder to facilitate the extraction process. Rhizome powder was extracted with 96% ethanol solvent for 3 days. The extraction results,

concentrated using a rotary evaporator at a temperature of $50-60^{\circ}$ C. Then, the extract was dried over water bath at $50-60^{\circ}$ C.

Screening of Phytochemistry Content

The dried extract, used for phytochemical screening and examination of the quality of the extract. Phytochemical screening was performed to determine the content of active compounds in the extracts including flavonoids, tannins, saponins, triterpenoids, and steroids. Extract quality checks include moisture content, total ash content, water soluble extract, and ethanol-soluble extract content.

Animals

This study was conducted on male Wistar rats, age 2 months, body weight 160-200 g. Animals adapted to laboratory conditions in enclosures with a light dark cycle of 12 hours, receiving standard food, drinking water ad libitum. The test protocol has been approved by the animal ethics committee, medical faculty, Padjadjaran University, Bandung, Indonesia (315 / UN6.C1.3.2 / KEPK / PN / 2016).

Study Design

A total of 30 male Wistar rats were divided randomly into 6 groups each 5 rats: group 1 (received drug carrier), group 2 (received drug carrier), group 3 (received simvastatin dose 0.9 mg / kg), group 4-6 received extract dose 30, 50, and 100 mg / kg. All groups (except group 2) received 25% fructose in drinking water for 21 days.

On the 22nd day, blood is taken to get serum. Serum is used for measurement of total cholesterol, LDL, HDL, and triglyceride levels. Total cholesterol and triglycerides were determined enzymatically using a reagent kit Proline®, while measurement of HDL and LDL cholesterol levels using a reagent kit Sekisui®. The color change in the sample was measured using a MicroLab 300® at a wavelength of 340 nm.

HMG CoA reductase Inhibitory Assay

Furthermore, animals were sacrificed and isolated liver and cleaned in 0.9% NaCl solution. The cleansed liver, homogenized 20% in phosphate buffer pH 7.4 and centrifuged at 8000 rpm, then supernatant separated (homogenate). Homogenate is used to test the activity of the HMG CoA reductase enzyme.

A 50 mL of liver homogenates, 260 mL of phosphate buffer pH 7.4, 42 mL of 50 mM Na 2 EDTA solution, 60 mL solution of dithiothreitol (DTT) 100 mM (purchased from Sigma), 50 mL of 2.16 M KCl solution, and 12 mL of HMG CoA substrate solution (purchased from Sigma) was mixed using a vortex for 10 seconds and incubated at 37 ° C for 30 minutes. After an incubation period, then added 55 mL of 1,105 mM NADPH (purchased from Sigma) and remixed using a vortex for 10 seconds and then incubated for 10 minutes. The absorbance of the sample were measured by MicroLab 300® at a wavelength of 340 nm.

Data Analysis

The data obtained were analyzed statistically using SPSS. Data are expressed as mean \pm standard deviation. The p value <0.05 indicates a significant difference compare to control group.

RESULTS

The result of quality test of ethanol extract of *Zingiber cassumunar* rhizome was obtained ash content of 9,45%, 10% water content, water soluble content of 72,42% and soluble extract content of ethanol 87,69%. Water content in the extract should be no more than 10%, because water is a growth medium for microorganisms and also as a medium of enzymatic reactions that can decompose the active compound. The results of phytochemical screening, ethanol extract of *Zingiber cassumunar* rhizomes contain compounds: flavonoids, triterpenoids, tannins and steroids.

The results showed that administration of fructose 25% in drinking water for 21 days, increased total cholesterol, LDL, and triglyceride levels, and decreased HDL levels in the blood. Administration of *Zingiber cassumunar* ethanol extracts of 30, 50, and 100 mg / kg for 21 days can lower total cholesterol, LDL and triglyceride levels and increase HDL cholesterol levels (tables 1 and 2).

The group receiving the extract at a dose of 50 mg / kg showed a decrease in total cholesterol and LDL levels compared with those receiving 30 and 100 mg / kg extracts. However, the 30 and 100 mg / kg doses still showed a statistically significant decrease effect compared to group 1 (p <0.05).

The group receiving 30 and 50 mg / kg dose extracts showed significantly different triglyceride levels statistically (p <0.05) compared to group 1 (Table 1). The decreased triglyceride levels in the group receiving doses of 30 and 50 mg / kg were greater than those receiving the dose of 100 mg / kg. However, administration of the extract at a dose of 100 mg / kg can lower triglyceride levels by day 22 but not statistically significant (p> 0.05) compared to group 1 (table 1).

The dose of extracts of 50 and 100 mg / kg could increase statistically significant HDL levels (p < 0.05) compared to group 1 (table 2). The increase in HDL cholesterol levels at doses of 50 and 100 mg / kg was greater than the dose of 30 mg / kg. Administration of the extract at doses of 30 mg / kg can increase HDL cholesterol levels on day 22 but not statistically significant (p > 0.05) compared with group 1 (table 2).

The extract of doses of 30, 50 and 100 mg / kg for 21 days was able to inhibit the activity of the enzyme HMG CoA reductase in the liver. The inhibitory value of the enzyme activity of HMG CoA reductase was calculated as percent inhibition (table 4). The higher percent inhibition indicates the stronger the test drug inhibits the activity of the enzyme HMG CoA reductase.

		Treatment	Gloups	
Crown	TC ± SD (mg/dL)		$TG \pm SD (mg/dL)$	
Group	TO	T22	TO	T22
1	67.5 ± 7.7	121.9 ± 6.8	83.0 ± 5.5	182.9 ± 29.8
2	59.5 ± 5.5	68.3 ± 11.6*	83.0 ± 7.7	93.1 ± 11.5*
3	64.4 ± 10.1	$84.2 \pm 9.8^{*}$	83.2 ± 10.1	117.0 ± 12.4*
4	62.8 ± 5.7	100.0 ± 16.4*	84.6 ± 5.7	148.4 ± 9.5*
5	63.2 ± 5.8	86.9 ± 7.9*	83.7 ± 5.8	129.8 ± 16.0*
6	69.9 ± 7.02	93.3 ± 9.88*	83.6 ± 7.02	172.7 ± 24.0

Table 1. The Average Level of Total Cholesterol and Triglycerides Serum at Day 0 and Day 22 for All

Group 1: received drug carrier (CMC Na 1%) and 25% D-fructose Group 2: received drug carrier (CMC Na 1%)

Group 3: received simvastatin 0.9 mg/kg orally

Group 4: received extract dose of 30 mg/kg and 25% D-fructose Group 5: received extract dose of 50 mg/kg and 25% D-fructose Group 6: received extract dose of 100 mg/kg and 25% D-fructose *significant difference compared to group 1 (p<0.05)

Table 2. The Average Level of Cholesterol LDL and Cholesterol HDL Serum at Day 0 and Day 22 for All Treatment Groups

Crown	$LDL \pm SD (mg/dL)$		HDL ± SD (mg/dL)	
Group	TO	T22	T0	T22
1	16.2 ± 2.2	79.0 ± 8.6	19.2 ± 2.7	6.2 ± 1.7
2	19.9 ± 2.7	$\textbf{20,3} \pm \textbf{1.5*}$	21.2 ± 2.6	22.4 ± 3.9*
3	18.3 ± 2.0	$33.2 \pm 4.2*$	16.5 ± 13.0	21.5 ± 5.1*
4	17.0 ± 2.8	61.0 ± 3.4*	16.2 ± 13.0	10.3 ± 2.2
5	19.1 ± 3.0	$42.5\pm8.0^{\ast}$	16.1 ± 3.2	17.2 ± 4.3*
6	19.1 ± 4.0	$57.2 \pm 6.0 \ast$	16.6 ± 4.0	17.1 ± 4.3*

Group 1: received drug carrier (CMC Na 1%) and 25% D-fructose

Group 2: received drug carrier (CMC Na 1%)

Group 3: received simvastatin 0.9 mg/kg orally

Group 4: received extract dose of 30 mg/kg and 25% D-fructose Group 5: received extract dose of 50 mg/kg and 25% D-fructose Group 6: received extract dose of 100 mg/kg and 25% D-fructose *significant difference compared to group 1 (p<0.05)

Tabel 3. The Average Value of Absorbance in HMG CoA Reductase Assay for All Group of Treatment

Group	Absorbance ± Standard Deviation (SD)	
1	0.115 ± 0.037	
3	0.053 ± 0.031	
4	0.102 ± 0.005	
5	0.092 ± 0.020	
6	0.061 ± 0.020	

Group 1: received drug carrier (CMC Na 1%) and 25% D-fructose Group 3: received simvastatin 0.9 mg/kg orally

Group 4: received extract dose of 30 mg/kg and 25% D-fructose Group 5: received extract dose of 50 mg/kg and 25% D-fructose

Group 6: received extract dose of 100 mg/kg and 25% D-fructose

 Table 4. The Percentage Inhibition of HMG-CoA Reductase

 Activity for All Treatment Groups

Group	Percentage Inhibition ± Standard Deviation (%)	
3	53.9 ± 9.2	
4	11.3 ± 8.6	
5	20.0 ± 4.6	
6	47.0 ± 4.6	

Group 3: received simvastatin 0.9 mg/kg orally

Group 4: received extract dose of 30 mg/kg and 25% D-fructose Group 5: received extract dose of 50 mg/kg and 25% D-fructose Group 6: received extract dose of 100 mg/kg and 25% D-fructose

DISCUSSION

Study of the anti-hyperlipidemia activity of ethanolic extract of Zingiber cassumunar Roxb, aimed to test the ability of extract in improving lipid profile in animal models of fructose-induced hyperlipidemia. A diet of 25% fructose in drinking water for 21 days (with standard food) was able to raise total cholesterol and triglyceride levels more than twice their normal value, increasing LDL levels about 5 times their normal value, and being able to lower HDL levels 3 times its normal value. These results are in line with research conducted by R. Borate 2011, that the fructose diet increases total cholesterol, triglyceride, and LDL levels and decreases HDL [9]. Other studies also reported a high-fructose diet significantly increased the concentration of triglycerides and LDL cholesterol [10]. Studies on the women population in the US, reported that the fructose diet affects cardio metabolic biomarkers including lowering HDL cholesterol [11].

A 25% fructose diet in drinking water for 21 days can be used to make animal models of hyperlipidemia especially for studies that test the effects of the drug on LDL cholesterol levels. It is known that elevated levels of LDL cholesterol play a role in the process of atherosclerosis is the formation of foam cells in blood vessel endothelial cells that cause disruption of blood flow characterized by decreased elasticity of blood vessels (stiffness of blood vessels) [12].

Previous research has reported that a 25% fructose diet in drinking water in male Wistar rats can increase systolic and diastolic blood pressure and increase heart rate [13]. So a 25% fructose diet in 21 days can be used as an animal model with cardiovascular disorders. The mechanism of cardiovascular disorders is thought to occur through increased risk of atherosclerosis (elevated levels of oxidized LDL) [14], stiffness of blood vessels (increased systolic and diastolic blood pressure), and increased risk of heart failure (increase in heart rate) [15].

Although elevated levels of total cholesterol and a decrease in HDL cholesterol may increase the risk of cardiovascular disease, an increase in LDL cholesterol has been shown to have a very strong association with the risk of cardiovascular atherosclerosis [16]. Therefore, lowering LDL cholesterol levels is still the main therapeutic target for reducing the risk of cardiovascular disease [17]. Previous studies have shown that the risk of cardiovascular atherosclerosis occurs in a population of patients who do not achieve target LDL cholesterol levels [18].

Statins have been shown to have good ability to lower LDL cholesterol levels [19]. Currently statins are still the main drug of choice for hyperlipidemia. The use of statins has been shown to decrease the occurrence of heart attacks and increase life expectancy in patients at high risk of cardiovascular attacks [20].

Atherosclerosis occurs through a complex process and a long time. The progression of atherosclerosis is influenced by various factors including genetic, risk factors and patient lifestyle [21]. However, heart attacks due to atherosclerotic plaque occur suddenly. Therefore, ideally, LDL-lowering cholesterol drugs are given early on before the formation of atherosclerotic plaque or early stage of atherosclerosis that can still be repaired. However, the difficulty of making animal models of atherosclerosis, as well as the many factors that affect the occurrence of atherosclerosis leads to limited studies of hyperlipidemia drugs. This poses a challenge for the study of non-invasive methods for early detection of atherosclerotic plaque, which begins with decreased elasticity of blood vessels [20].

However, the use of long-term statins requires monitoring of liver function and rhabdomyolysis side effects. This increases the risk of patient non-adherence to the therapy so that the patient does not achieve the therapeutic target [22]. This, the challenge as well as the opportunity of herbal medicine as a source of new drugs to lower cholesterol, especially LDL cholesterol.

Based on the results of this study has proven that ethanolic extract of *Zingiber cassumunar* can improve lipid profile in animal models of fructose-induced hyperlipidemia. The dose extract of 50 mg / kg was able to lower LDL -46% cholesterol, triglyceride -29%, total cholesterol -29% and increase HDL cholesterol 3 times (177%). The results of this study, reinforce previous research, that ethanolic extract of *Zingiber cassumunar* can decrease heart rate in animal models of 25% fructose induced arrhythmias in drinking water, thereby reducing the risk of heart failure [13].

Our study was able to demonstrate the effect of *Zingiber* cassumunar in increasing HDL cholesterol by 3-fold. This is an opportunity to develop the potential benefits of *Zingiber cassumunar* very strongly in increasing HDL cholesterol, in addition to lowering LDL cholesterol. Previous studies have reported that HDL cholesterol plays a role in the regression of atherosclerotic plaque by decreasing inflammatory factors [23]. Therefore, increasing HDL cholesterol levels are a potential therapeutic target as atheroprotection in patients with high cardiovascular risk [24].

A decrease in LDL cholesterol can occur due to inhibition of HMG CoA reductase in the liver. This enzyme plays an important role in regulating endogenous cholesterol synthesis. The body's need for cholesterol is largely supplied through endogenous synthesis in the liver. However, the presence of genetic abnormalities in the number of LDL receptors, hyperactivity of HMG CoA reductase enzyme, and the effect of dietary excess cholesterol, causes hyperlipidemia.

Our results showed that ethanolic extract of *Zingiber cassumunar* have the ability to inhibit the activity of the HMG-CoA reducing enzyme in the liver with increased inhibition percentage in accordance with increasing the dose of the extract. The percentage of inhibition obtained is used to calculate the IC50 value of the linear regression equation. The dose of *Zingiber cassumunar* extract which can inhibit 50% of HMg CoA reductase enzyme activity was 106 mg / kg.

HMG-CoA reductase is the rate-limiting enzyme of cholesterol synthesis in the liver. This enzyme as a catalyst in the cholesterol biosynthesis reaction. HMG CoA reductase is the target of statin drugs, as a competitive inhibitor of HMG CoA reductase enzyme activity. Statins bind to the active side of the HMG CoA reductase, thus inhibiting substrate access (HMG CoA) against the active site. While NADPH, as the second substrate of the enzyme, is not occupied by inhibitor molecules. As the effect of the reaction is decreased cholesterol synthesis in the liver so that cholesterol levels in blood decreases [25].

The results of phytochemical screening of ethanolic extract of *Zingiber cassumunar* showed the presence of secondary metabolite compounds including flavonoids and tannins. These results are in line with studies reporting that *Zingiber cassumunar* contains various bioactive compounds that have various pharmacological activities such as antiinflammatory, antimicrobial, anticancer, antioxidant, antimalarial [26]. Flavonoids and tannins in *Zingiber cassumunar* are thought to be active compounds that act as inhibitors of HMG CoA reductase enzyme activity. However, it requires further research to prove it. Other studies have reported that the active compound in the ethanolic extract of *Zingiber cassumunar* has an antiinflammatory effect [26].

CONCLUSION

The results of this study, it can be concluded that *Zingiber* cassumunar Roxb ethanol extract can show the activity of anti-hyperlipidemia in animal models of fructose-induced hyperlipidemia. One of the mechanisms of action of *Zingiber cassumunar* ethanol extract in lowering cholesterol through inhibition of HMG CoA reductase enzyme, an enzyme that plays an important role in the synthesis of endogenous cholesterol in the liver.

REFERENCES

- Dennison-Himmelfarb C, Handler J, Lackland DT, LeFevre ML, MacKenzie TD, Ogedegbe O, et al. 2014 Evidence-Based Guideline for the Management of High Blood Pressure in Adults Report From the Panel Members Appointed to the Eighth Joint National Committee (JNC 8). JAMA. 2014;311(5):507–20.
- 2. Mendis S, Puska P, Norrving B, Organization WH. Global atlas on cardiovascular disease prevention and control. Geneva: World

Health Organization; 2011.

- Kementerian Kesehatan I. RISKESDAS 2013. Jakarta Kementeri Kesehat RI. 2013;
- Jena M, Jena J, Biswal SB, Mishra S, Pal A. Effect of Eclipta alba on fructose induced hypertension in albino rats. Int J Pharm Pharm Sci. 2013;5(3):281–5.
- Laughlin MR, Bantle JP, Havel PJ, Parks E, Klurfeld DM, Teff K, et al. Clinical research strategies for fructose metabolism. Adv Nutr. 2014;5(3):248–59.
- 6. Monteiro R, Azevedo I. Chronic inflammation in obesity and the metabolic syndrome. Mediators Inflamm. 2010;2010:220-220.
- Mukhtar RYA, Reckless JPD. Statin-induced myositis: a commonly encountered or rare side effect? Curr Opin Lipidol. 2005;16(6):640– 7.
- Singh C, Manglembi N, Swapana N, Chanu S. Ethnobotany, phytochemistry and pharmacology of Zingiber cassumunar Roxb.(Zingiberaceae). J Pharmacogn Phytochem. 2015;4(1).
- Borate AR, Suralkar AA, Birje SS, Malusare P V, Bangale PA. Antihyperlipidemic effect of protocatechuic acid in fructose induced hyperlipidemia in rats. Int J Pharma Bio Sci. 2011;2(4):456–60.
- Schaefer EJ, Gleason JA, Dansinger ML. Dietary fructose and glucose differentially affect lipid and glucose homeostasis. J Nutr. 2009;139(6):1257S–1262S.
- Yu Z, Ley SH, Sun Q, Hu FB, Malik VS. Cross-sectional association between sugar-sweetened beverage intake and cardiometabolic biomarkers in US women. Br J Nutr. 2018;119(5):570–80.
- Badimon L, Vilahur G. LDL-cholesterol versus HDL-cholesterol in the atherosclerotic plaque: inflammatory resolution versus thrombotic chaos. Ann N Y Acad Sci. 2012;1254(1):18–32.
- Hasimun P, Sulaeman A, Sutrisno E, Putra HM. POTENTIAL DIURETIC ACTIVITY FROM ZINGIBERACEAE FAMILY AND ITS EFFECT ON THE LEVELS OF POTASSIUM AND SODIUM IONS IN THE URINE. World J Pharm Pharm Sci [Internet]. 2018;7(3):226–36.
- 14. Mitra S, Goyal T, Mehta JL. Oxidized LDL, LOX-1 and atherosclerosis. Cardiovasc Drugs Ther. 2011;25(5):419.

- Wagenseil JE, Mecham RP. Elastin in large artery stiffness and hypertension. J Cardiovasc Transl Res. 2012;5(3):264–73.
- Ference BA. Causal Effect of Lipids and Lipoproteins on Atherosclerosis: Lessons from Genomic Studies. Cardiol Clin. 2018;36(2):203–11.
- Hadjiphilippou S, Ray KK. Lipids and Lipoproteins in Risk Prediction. Cardiol Clin. 2018;36(2):213–20.
- Zhang M, Deng Q, Wang L, Huang Z, Zhou M, Li Y, et al. Prevalence of dyslipidemia and achievement of low-density lipoprotein cholesterol targets in Chinese adults: A nationally representative survey of 163,641 adults. Int J Cardiol. 2018;260:196–203.
- Würtz P, Wang Q, Soininen P, Kangas AJ, Fatemifar G, Tynkkynen T, et al. Metabolomic profiling of statin use and genetic inhibition of HMG-CoA reductase. J Am Coll Cardiol. 2016;67(10):1200–10.
- 20. Goldstein JL, Brown MS. A century of cholesterol and coronaries: from plaques to genes to statins. Cell. 2015;161(1):161–72.
- Rafieian-Kopaei M, Setorki M, Doudi M, Baradaran A, Nasri H. Atherosclerosis: process, indicators, risk factors and new hopes. Int J Prev Med. 2014;5(8):927.
- Thompson PD, Panza G, Zaleski A, Taylor B. Statin-associated side effects. J Am Coll Cardiol. 2016;67(20):2395–410.
- Feig JE, Rong JX, Shamir R, Sanson M, Vengrenyuk Y, Liu J, et al. HDL promotes rapid atherosclerosis regression in mice and alters inflammatory properties of plaque monocyte-derived cells. Proc Natl Acad Sci. 2011;108(17):7166–71.
- Rader DJ, Hovingh GK. HDL and cardiovascular disease. Lancet [Internet]. 2014;384(9943):618–25.
- Istvan ES. Structural mechanism for statin inhibition of 3-hydroxy-3methylglutaryl coenzyme A reductase. Am Heart J. 2002;144(6):S27–32.
- Khemawoot P, Hunsakunachai N, Anukunwithaya T, Bangphumi K, Ongpipattanakul B, Jiratchariyakul W, et al. Pharmacokinetics of compound D, the major bioactive component of Zingiber cassumunar, in rats. Planta Med. 2016;82(13):1186–91.