

DAFTAR PUSTAKA

- Aditya, N. P., Macedo, A. S., Doktorovova, S., Souto, E. B., Hoon, S., Chang, P., & Ko, S. (2014). Development and evaluation of lipid nanocarriers for quercetin delivery: a comparative study of solid lipid nanoparticles (SLN), nanostructured lipid carriers (NLC), and lipid nanoemulsions (LNE). *LWT - Food Science and Technology*. <https://doi.org/10.1016/j.lwt.2014.04.058>
- Amanina, N. I. K., Abu, F., Sahudin, S., Hussain, Z., & Hussain, M. (2018). SELF-ASSEMBLED CHITOSAN NANOPARTICLES FOR PERCUTANEOUS DELIVERY OF CAFFEINE : PREPARATION , CHARACTERIZATION AND IN VITRO RELEASE STUDIES, *10*(4).
- Ascenso, A., & Marques, H. C. (2009). Acne in the Adult. *Mini-Reviews in Medicinal Chemistry*, *9*(1), 1–10.
- Budiputra, dewa ken. (2013). *ENGEMBANGAN FORMULA DAN KARAKTERISASI DALAM BENTUK GEL UNTUK RUTE TRANSDERMAL DEWA KEN BUDIPUTRA NIM : 20711017 (Program Studi Farmasi) INSTITUT TEKNOLOGI BANDUNG*.
- Casey, G. (2002). Physiology of the skin. *Nursing Standard (Royal College of Nursing (Great ...)*, *47–51*.
<https://doi.org/10.1016/B978-1-4831-7818-9.50015-0>
- Charoenputtakhun, P., Opanasopit, P., Rojanarata, T., & Ngawhirunpat, T. (2013). All-trans retinoic acid-loaded lipid nanoparticles as a transdermal drug delivery carrier, *7450*(66), 1–9. <https://doi.org/10.3109/10837450.2013.763261>
- Chen, Y., Zhou, L., Yuan, L., Zhang, Z. hai, Liu, X., & Wu, Q. (2012). Formulation, characterization, and evaluation of in vitro skin permeation and in vivo pharmacodynamics of surface-charged tripterine-loaded nanostructured lipid carriers. *International Journal of Nanomedicine*, *7*, 3023–3033.

<https://doi.org/10.2147/IJN.S32476>

- Cheng-ying, S., Ling, D. A. I., Bao-de, S., Xu, Z., Jin-xia, B. A. I., He, X. U., ... Hai-long, Y. (2015). Nanostructured lipid carrier based topical gel of Ganoderma Triterpenoids for frostbite treatment, *13*(6), 454–460. [https://doi.org/10.1016/S1875-5364\(15\)30039-X](https://doi.org/10.1016/S1875-5364(15)30039-X)
- Danaei, M., Dehghankhold, M., Ataei, S., Hasanzadeh Davarani, F., Javanmard, R., Dokhani, A., ... Mozafari, M. R. (2018). Impact of particle size and polydispersity index on the clinical applications of lipidic nanocarrier systems. *Pharmaceutics*, *10*(2), 1–17. <https://doi.org/10.3390/pharmaceutics10020057>
- Donalisio, M., Leone, F., Lembo, D., & Cavalli, R. (2018). Acyclovir-Loaded Chitosan Nanospheres from Nano-Emulsion Templating for the Topical Treatment of Herpesviruses Infections, *1*, 1–12. <https://doi.org/10.3390/pharmaceutics10020046>
- Gao, Y., Han, K., Wang, Q., Hu, Z., Liu, Q., Liu, L., & Zeng, K. (2018). Development of podophyllotoxin-loaded nanostructured lipid carriers for the treatment of condyloma acuminatum. *Molecular Medicine Reports*, 6506–6514. <https://doi.org/10.3892/mmr.2018.8696>
- Garg, T. (2014). Current nanotechnological approaches for an effective delivery of bio-active drug molecules in the treatment of acne. *Artificial Cells, Nanomedicine and Biotechnology*, *44*(1), 98–105. <https://doi.org/10.3109/21691401.2014.916715>
- Gonullu, U., Uner, M., Yuner, G., Karaman, ecem fatma, & Aydogmus, Z. (2015). Formulasi dan karakterisasi nanopartikel lipid padat , berstruktur nano operator lipid dan nanoemulsion dari lornoxicam, *65*, 1–13.
- Hang, X. Z., An, W. P., An, L. G., Hu, C. Z., An, Y. G., & Ie, S. N. (2008). Preparation of a Dispersible PEGylate Nanostructured Lipid Carriers (NLC) Loaded with 10-Hydroxycamptothecin

by Spray-Drying, 56(December), 1645–1650.

- Haryanto, isnaeni yudi, & Hendradi, E. (2017). Physical and chemical characteristics of meloxicam from nanostructured lipid carriers system using so *Asian Journal of Pharmaceutical and Clinical Research*, 10(2), 132–137.
<https://doi.org/10.22159/ajpcr.2017.v10i2.14806>
- Jain, A. K., Jain, A., Garg, N. K., Jain, A., Jain, S. A., Tyagi, R. K., & Agrawal, G. P. (2014). Ac ce p t e d t. In *Colloids and Surfaces B: Biointerfaces*. Elsevier B.V.
<https://doi.org/10.1016/j.colsurfb.2014.05.041>
- Kakadia, P., & Conway, B. (2015). *Lipid Nanoparticles for Dermal Drug Delivery*. *Current Pharmaceutical Design* (Vol. 21). Elsevier Inc.
<https://doi.org/10.2174/1381612821666150428143730>
- Kielhorn, J., & International Programme on Chemical Safety (Eds.). (2006). *Dermal absorption*. Geneva: WHO.
- Kim, B., Na, Y., Choi, J., Kim, I., Lee, E., Kim, S., ... Cho, C. (2017). The Improvement of Skin Whitening of Phenylethyl Resorcinol by Nanostructured Lipid Carriers. *Journal Nanomaterials*, 241(7), 1–17.
<https://doi.org/10.3390/nano7090241>
- Kovacevic, A., Savic, S., Vuleta, G., Müller, R. H., & Keck, C. M. (2011). Polyhydroxy surfactants for the formulation of lipid nanoparticles (SLN and NLC): Effects on size , physical stability and particle matrix structure. *International Journal of Pharmaceutics*, 406(1–2), 163–172.
<https://doi.org/10.1016/j.ijpharm.2010.12.036>
- Kurniawati, A. R. (2014). Kejadian Acne Vulgaris.
- Loo, M., & Press, D. (2013). Effect of compositions in nanostructured lipid carriers (NLC) on skin hydration and occlusion. *International Journal of Nanomedicine*, 13–22.

- López-García, R., & Ganem-Rondero, A. (2015). Solid Lipid Nanoparticles (SLN) and Nanostructured Lipid Carriers (NLC): Occlusive Effect and Penetration Enhancement Ability. *Journal of Cosmetics, Dermatological Sciences and Applications*, 05(02), 62–72.
<https://doi.org/10.4236/jcdsa.2015.52008>
- Madelina, W., & silistyaningsih. (2018). Farmaka Farmaka. *Farmaka*, 16(2), 105–117.
- Mandawgade, S. D., & Patravale, V. B. (2008). Development of SLNs from natural lipids : Application to topical delivery of tretinoin. *Pharmaceutical Nanotechnology*, 363, 132–138.
<https://doi.org/10.1016/j.ijpharm.2008.06.028>
- Mappamasing, F., Anwar, E., & Im, A. M. U. N. (2015). Formulasi , Karakterisasi dan Uji Penetrasi In Vitro Resveratrol Solid Lipid Nanopartikel dalam Krim Topikal (Formulation , Characterization and In Vitro Penetration Study of Resveratrol Solid Lipid Nanoparticles in Topical Cream). *Jurnal Ilmu Kefarmasian Indonesia*, 13(2), 137–144.
- Mehnert, W., & Mader, K. (2001). Solid lipid nanoparticles Production , characterization and applications, 47, 165–196.
- Müller, R. H., Petersen, R. D., Hommoss, A., & Pardeike, J. (2007). Nanostructured lipid carriers (NLC) in cosmetic dermal products. *Advanced Drug Delivery Reviews*, 59(6), 522–530.
<https://doi.org/10.1016/j.addr.2007.04.012>
- noviyanti, sintia dwi. (2018). pengembangan formula nanostructured lipid carrier (NLC) alfa-tokoferol.
- Pardeike, J., Hommoss, A., & Müller, R. H. (2009). Lipid nanoparticles (SLN , NLC) in cosmetic and pharmaceutical dermal products, 366, 170–184.
<https://doi.org/10.1016/j.ijpharm.2008.10.003>
- Phatsawee, J., Zoltán, F., & C, R. G. (2019). Ac ce pt us. In *Drug*

Development and Industrial Pharmacy (Vol. 0, p. 000). Taylor & Francis. <https://doi.org/10.1080/03639045.2019.1569023>

Prihapsara, F. (2018). Optimization and Antioxidant Activity Test Of SMEDDS Bay Leaf with Palm Kernell Oil as Oil Phase. *JPSCR : Journal of Pharmaceutical Science and Clinical Research*, 2(02), 102. <https://doi.org/10.20961/jpscr.v2i02.14390>

Rullah, K., Lucida, H., & Hamidi, D. (2012). Pengembangan potensi rhodomyrtone sebagai bahan aktif sediaan topikal. *Jurnal Farmasi Indonesia*, 6(1), 7–12.

Sally, S., H, I. R. A., M, H. R., & D, M. N. (2017). Nanostructured Lipid Carriers Loaded with Simvastatin : Effect of PEG / glycerides on characterization , stabilit Nanostructured lipid carriers loaded with simvastatin : effect of PEG / glycerides on characterization , stability , cellular uptake. *Drug Development and Industrial Pharmacy*, 0(0), 000. <https://doi.org/10.1080/03639045.2017.1293681>

Sampelan, M. G., & Kundre, D. P. R. M. (2017). Hubungan timbulnya acne vulgaris dengan tingkat kecemasan pada remaja di SMP N 1 Likupang Timur. *E-Journal Keperawatan (e-Kp)*, 5(1), 2.

Shah, M. R., Imran, M., & Ullah, S. (2017). *Lipid-Based Nanocarriers for Drug Delivery and Diagnosis. Lipid-Based Nanocarriers for Drug Delivery and Diagnosis*. <https://doi.org/https://doi.org/10.1016/B978-0-323-52729-3.00005-6>

Shalita, A., Weiss, J. S., Chalker, D. K., Ellis, C. N., & Arbor, A. (1996). A comparison of the efficacy and safety of adapalene gel 0 . 1 % and tretinoin gel 0 . 025 % in the treatment of acne vulgaris : A multicenter trial. *Journal of the American Academy of Dermatology*, 34, 482–485.

Shamma, rehab nabil, & Aburahma, mona hassan. (2014).

Follicular delivery of spironolactone via nanostructured lipid carriers for management of alopecia. *International Journal of Nanomedicine*, 5449–5460.

Souza, L. G., Silva, E. J., Martins, A. L. L., Mota, M. F., Braga, R. C., Lima, E. M., ... Marreto, R. N. (2011). European Journal of Pharmaceutics and Biopharmaceutics Development of topotecan loaded lipid nanoparticles for chemical stabilization and prolonged release. *European Journal of Pharmaceutics and Biopharmaceutics*, 79(1), 189–196.
<https://doi.org/10.1016/j.ejpb.2011.02.012>

Tamjidi, F., Shahedi, M., Varshosaz, J., & Nasirpour, A. (2013). *Nanostructured lipid carriers (NLC): A potential delivery system for bioactive food molecules. Innovative Food Science and Emerging Technologies* (Vol. 19). Elsevier B.V.
<https://doi.org/10.1016/j.ifset.2013.03.002>

Theresia, M. (2013). Acne vulgaris. *Continuing Medical Education*, 40(8), 58. <https://doi.org/10.1007/s00105-010-1918-7>

Tofani, R. P., Sumirtapura, Y. C., & Darijanto, S. T. (2016). Formulation , Characterisation , and in Vitro Skin Diffusion of Nanostructured Lipid Carriers for Deoxyarbutin Compared to a Nanoemulsion and Conventional Cream. In *scientia pharmaceutica* (pp. 634–645).
<https://doi.org/10.3390/scipharm84040634>

Upreti, T. (2017). Nanostructured Lipid Carrier System for the Treatment for Skin Disease-A Review. *JSM Nanotechnology & Nanomedicine*, 5(3), 1059. Retrieved from <https://www.jscimedcentral.com/Nanotechnology/nanotechnology-5-1059.pdf>

Veerapur, V. P. (2014). *International Journal of Biopharmaceutics*, 5(3), 218–224.

Wells, barbara g, & Dipiro, joseph t. (2015). *Pharmacotherapy Handbook*, 135.

Wolf, M., Klang, V., Stojcic, T., Fuchs, C., Wolzt, M., & Valenta, C. (2018). NLC versus nanoemulsions: Effect on physiological skin parameters during regular in vivo application and impact on drug penetration. *International Journal of Pharmaceutics*, 549(1–2), 343–351.
<https://doi.org/10.1016/j.ijpharm.2018.08.007>

National Center for Biotechnology Information. PubChem Database. Adapalene, CID=60164, <https://pubchem.ncbi.nlm.nih.gov/compound/Adapalene> (accessed on July 1, 2019)

National Center for Biotechnology Information. PubChem Database. Glyceryl monostearate, CID=24699, <https://pubchem.ncbi.nlm.nih.gov/compound/Glyceryl-monostearate> (accessed on July 1, 2019)