

DAFTAR PUSTAKA

- Amelinda, E., Widarta, I. W. R., & Darmayanti, L. P. T. (2018). PENGARUH WAKTU MASERASI TERHADAP AKTIVITAS ANTIOKSIDAN EKSTRAK RIMPANG TEMULAWAK (*Curcuma xanthorrhiza* Roxb.). *Jurnal Ilmu Dan Teknologi Pangan (ITEPA)*, 7(4), 165. <https://doi.org/10.24843/itepa.2018.v07.i04.p03>
- Atun, S., Aznam, N., Arianingrum, R., Senam, Naila, B. I. A., Lestari, A., & Purnamaningsih, N. A. (2020). Characterization of curcuminoid from curcuma xanthorrhiza and its activity test as antioxidant and antibacterial. *Molekul*, 15(2), 79–87. <https://doi.org/10.20884/1.jm.2020.15.2.540>
- Hayani, E. (2006). Analisis Kandungan Kimia Rimpang Temulawak. *Temu Teknis Nasional Tenaga Fungsional Pertanian.*, 309–312.
- Jain, P. K., & Agrawal, R. K. (2008). Antioxidant and Free Radical Scavenging Properties of Developed Mono-and Polyherbal Formulations. *Asian J. Exp. Sci*, 22(3), 213–220.
- jun, fu, j. hong. (2003). Comparison of antioxidant activities of isoflavones from kudzu root. *Jurnal of Food Science*, 68(6), 2117–2122. <https://doi.org/10.1111/j.1365-2621.2003.tb07029.x>
- Nurcholis, W., & Bintang, M. (2017). Perbandingan Aktivitas Antioksidan dan Kandungan Fenolik Temulawak dan Temu Ireng. *Jurnal Jamu Indonesia*, 2(1), 25–29. <https://doi.org/10.29244/jji.v2i1.27>
- Qonitah, F. (2020). Aktivitas Antioksidan Krim “X” Dengan Metode DPPH ((1,1-Diphenyl-2-Picryl-Hidrazyl). *Pharmed: Journal of Pharmaceutical Science and Medical Research*, 3(2), 68. <https://doi.org/10.25273/pharmed.v3i2.7604>
- Rahmawati, R., Muflihunna, A., & Sarif, L. M. (2016). ANALISIS AKTIVITAS ANTIOKSIDAN PRODUK SIRUP BUAH MENGKUDU (*Morinda citrifolia* L.) DENGAN METODE DPPH. *Jurnal Fitofarmaka Indonesia*, 2(2), 97–101. <https://doi.org/10.33096/jffi.v2i2.177>
- Rosidi, A. (2020). The difference of Curcumin and Antioxidant activity in Curcuma

- Xanthorrhiza at different regions. *Journal of Advanced Pharmacy Education & Research*, 10(1), 14–18. www.japer.in
- Rosidi, A., Khomsan, A., Setiawan, B., & Briawan, D. (2004). Potensi temulawak (c. *Potensi Temulawak*, 1995.
- Septiana, D., Masruriati, E., & Fajaryanti, N. (2019). Uji Stabilitas Fisik Krim Ekstrak Etanol Temulawak Dengan Perbedaan Konsentrasi Asam Stearat. *Dalam: Jurnal Farmasetis, Sekolah Tinggi Ilmu Kesehatan, Kendal.*, 9(1), 7–14.
- Trifena. (2012). Analisis Uji in Vitro dan in Vivo Ekstrak Kombinasi Kulit Manggis (*Garcinia mangostana* L.) dan Pegagan (*Centella asiatica* L.) sebagai Krim Antioksidan. *Fmipa Ui*, 1–117.
- Vinchure, B. D., Kokate, R., & Patil, S. P. (2017). Formulation and evaluation of cream containing curcumin and lycopene for the antioxidant and antibacterial activity. *EUROPEAN JOURNAL OF PHARMACEUTICAL and Medical Research*, 4(5), 394–399.
- Prana, MS. 2008. The biology of temulawak (*Curcuma xanthorrhiza* Roxb.). Bogor (ID) : Biopharmaca Research Center Bogor Agricultural University. Hal. 151-156.
- Murray R. K., Granner D.K., Rodwell V.W., 2009. Biokimia Harper, (Andri Hartono)..Edisi 27.Penerbit Buku Kedokteran, EGC. Jakarta.
- [WHO] World Health Organization. 1999. Monograph on selected medicinal plant. Vol 1. Jenewa: WHO
- Sidik, Mulyono MW, Muhtadi A. 1992. Temulawak (*Curcuma xanthorrhiza* Roxb). Jakarta (ID) : Yayasan Pengembangan Obat Bahan Alam Phytomedica
- Fahn, A. 1991. Anatomi Tumbuhan edisi ketiga. Gadjah Mada University Press. Yogyakarta.
- Sayuti, MS, P., & Yenrina, MSi, D. (2015). Antioksidan Alami dan Sintetik. Padang: Andalas University Press.

Jain, P. K., & Agrawal, R. K. (2008). Antioxidant and Free Radical Scavenging Properties of Developed Mono- and Polyherbal Formulations. *Asian J. Exp. Sci*, 22(3), 213–220.