

DAFTAR PUSTAKA

- Juarta T, dr.J, Arifah M, Laksamana R 2024. “PEDOMAN PENATAKLAKSANAAN ANESTESI REGIONAL”
- Valiani C, Sugiarti I, Kartadarma S, Widayanti L, Ramadani A. 2025. “HUBUNGAN HIPOTENSI INTRAOPERATIF TERHADAP KEJADIAN POST OPERATIF NAUSEA AND VOMITING (PONV) PADA PASIEN PASCA SPINAL ANESTESI SEKSIO SEKSAREA Cici Valiani, Ina Sugiarti, Supriyatni Kartadarma, Lastri Widayanti, Anisa Ramadani.” 15: 59–66. <http://journal.stikeskendal.ac.id/index.php/PSKM>.
- Farida, F. & Handajanti, D. O. (2024). Pengaruh mobilisasi dini terhadap tingkat nyeri luka pada ibu post *sectio caesarea* metode ERACS di RSI Darus Syifa' Surabaya. *Indonesian Journal of Midwifery Today*, 4(1), 1–7.
- Gisbella C, Susanto A, Lintang, and Program. 2024. “GAMBARAN KEJADIAN POST OPERATIVE NAUSEA AND VOMITING (PONV) PADA PASIEN PASCA OPERASI SECTIO CAESAREA DI RSUD CILACAP”2(September):72133.[file:///C:/Users/ACER/Downloads/scientica+GAMBARAN+KEJADIAN+POST+OPERATIVE+NAUSEA+AND+VOMITING+\(PONV\)+PADA+PASIEN+PASCA+OPERASI+SECTIO+CAESAREA+DI+RSUD+CILACAP.pdf](file:///C:/Users/ACER/Downloads/scientica+GAMBARAN+KEJADIAN+POST+OPERATIVE+NAUSEA+AND+VOMITING+(PONV)+PADA+PASIEN+PASCA+OPERASI+SECTIO+CAESAREA+DI+RSUD+CILACAP.pdf).
- Akyol, D., Çelik, M., Ay, N., & Yıldız, G.Ö. (2024). *The effect of spinal needle type on post-dural puncture headache in spinal anesthesia: Prospective randomized study. Eurasian Journal of Medicine*, 56(1), 42–46.
- Manuputty, B. T. (2022). Gambaran nyeri punggung pada pasien pasca spinal anastesi di rsu negara.https://repository.itekesbali.ac.id/medias/journal/Beatrice_Tamariska_Manuputty.pdf

- Shahzada, G., Andrabi, S. W., Shaiq, H., & Malla, S. M. (2018). *Spinal anesthesia at L2–3 vs L3–4 levels: comparison of onset of block, maximum level of block and hemodynamic response in obstetric patients undergoing caesarian section. Journal of Medical Science and Clinical Research*, 6(4), 4778–4783.
- Olawin & M Das, (2023) Di ruang subarachnoid antara vertebra lumbal 2-3, 3-4, atau 4-5. (2021). *Pharmacognosy Magazine*, 75(17), 399–405.
- Arlena, N., Novitasari, D., & Burhan, A. (2024). *Differences in the time to achieve Bromage Score 2 in post-spinal anesthesia caesarean section patients between the use of conventional methods and ERACS. Indonesian Journal of Global Health Research*, 7(1), 389–398.
- Vishwanath, D., Kumar, M., & Bhat, S. (2021). *Effect of injection rate of hyperbaric bupivacaine on sensory block characteristics in spinal anesthesia: A randomized controlled trial. Clinical and Experimental Obstetrics & Gynecology*, 49(7), 152–157.
- Katz, J., & Matjasko, L. (2002). *The role of the anesthetist in preventing intraoperative nausea and vomiting. British Journal of Anaesthesia*, 89(5), 729-738
- Sakai, T., Aoyama, K., & Kuroda, M. (2013). *Comparison of spinal anesthesia at L2-L3 versus L3-L4: A randomized controlled trial. Anesthesia and Analgesia*, 116(2), 391-397.
- Zhou, J., & Zhang, C. (2017). The impact of spinal anesthesia injection sites on sensory and motor block characteristics. *Journal of Anesthesia & Clinical Research*, 8(12), 1006-1010.
- Ruli Herman Sitanggang, Siti Harniati, I. S. R. (2015). Perbandingan Pengaruh Penambahan Fentanil 25 Mg dengan Midazolam 2 Mg pada Bupivakain Hiperbarik 10 Mg Terhadap Lama Analgesia dan Tekanan Darah pada Seksio Sesaria. *Departemen Anestesiologi Dan Terapi Intensif Fakultas*

Kedokteran Universitas Padjadjaran RSUP. Dr. Hasan Sadikin-Bandung, 33, 252–261.

Triola, M. F. (2018). *Elementary statistics* (13th ed.). Pearson.

Nandaswari, N. M. N., Parami, P., Suranadi, I. W., & Senapathi, T. G. A. (2024). *Incidence of hypotension and nausea vomiting in subarachnoid block anesthesia technique in cesarean surgery*. *E-Jurnal Medika Udayana*, 13(3).

Chahar, P., et al. (2022). *Comparative evaluation of three techniques for paramedian spinal anesthesia: Real-time ultrasound-guided, pre-procedural ultrasound-assisted, versus landmark-based*. *Indian Journal of Anaesthesia*, 66(Suppl 2), S102–S107.

Xiao, F., et al. (2025). *Effect of injectate temperature on ED₅₀ of intrathecal hyperbaric bupivacaine for caesarean section: A randomized double-blind trial*. *Anesthesia & Analgesia*. Advance online publication

Asghar, S., et al. (2025). *Comparison of spinal anesthesia at L3–4 versus L4–5 in obstetric patients undergoing caesarean section*. *The Professional Medical Journal*. Advance online publication.

Pavičić Šarić, J., Mikulandra, S., Gustin, D., & Matasic, H. (2012). *Spinal anesthesia at the L2–3 and L3–4 levels: Comparison of analgesia and hemodynamic response*. *Collegium Antropologicum*, 36(1), 151–156.

Shin, D. W., Kim, Y., Hong, B., Yoon, S.-H., Lim, C. S., & Youn, S. (2019). *Effect of fentanyl on nausea and vomiting in cesarean section under spinal anesthesia: A randomized controlled study*. *Journal of International Medical Research*, 47(10), 4798–4807.

Dyer, R. A., Reed, A. R., van Dyk, D., Arcache, M. J., & Lombard, C. J. (2021). *Prevention and management of spinal hypotension during cesarean section*. *International Journal of Obstetric Anesthesia*, 47, 103207.

Chekol, W. B., Tadele, A. M., & Yimer, H. T. (2021). *Prevalence and associated factors of intraoperative nausea and vomiting during cesarean section under spinal anesthesia at public hospitals in Southern Ethiopia. Anesthesiology Research and Practice*, 2021, Article ID 3746359.